



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/53, 15/82, 9/02, 5/00, A01H 5/00, 5/10</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/19493</b> <b>(43) International Publication Date:</b> 22 April 1999 (22.04.99)
<b>(21) International Application Number:</b> PCT/US98/20807 <b>(22) International Filing Date:</b> 5 October 1998 (05.10.98) <b>(30) Priority Data:</b> 08/948,564 10 October 1997 (10.10.97) US <b>(71) Applicant (for all designated States except US):</b> NORTH CAROLINA STATE UNIVERSITY [US/US]; 103 Bynum Hall, CB #7003, Raleigh, NC 27695-7003 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SIMINSZKY, Balazs [HU/US]; 708 F Chapel Hill Drive, Raleigh, NC 27606 (US). DEWEY, Ralph, E. [US/US]; 9432 Cartersville Court, Raleigh, NC 27613 (US). CORBIN, Frederick, T. [US/US]; 4508 Lead Mine Road, Raleigh, NC 27612 (US). <b>(74) Agents:</b> BENNETT, Virginia, C. et al.; Myers, Bigel, Sibley & Sajovec, P.A., P.O. Box 37428, Raleigh, NC 27627 (US).		<b>(81) Designated States:</b> AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> CYTOCHROME P-450 CONSTRUCTS AND METHODS OF PRODUCING HERBICIDE-RESISTANT TRANSGENIC PLANTS  <b>(57) Abstract</b>  DNA sequence encoding novel cytochrome P-450 molecules are provided. The use of DNA constructs containing such molecules to transform plants is described, as are transgenic plants exhibiting increased resistance to phenylurea herbicides. Methods of using such DNA constructs and transformed plants are provided.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## NOVEL CYTOCHROME P-450 CONSTRUCTS AND METHODS OF PRODUCING HERBICIDE-RESISTANT TRANSGENIC PLANTS

### Field of the Invention

The present invention relates to DNA encoding novel cytochrome P-450 molecules, and the transformation of cells with such DNA. These DNA sequences may be used in methods of producing plants with an altered ability to  
5 metabolize chemical compounds, such as phenylurea herbicides.

### Background of the Invention

Cytochrome P-450 (P-450) monooxygenases are ubiquitous hemoproteins present in microorganisms, plants and animals. Comprised of a large and diverse  
10 group of isozymes, P-450s mediate a great array of oxidative reactions using a wide range of compounds as substrates, and including biosynthetic processes such as phenylpropanoid, fatty acid, and terpenoid biosynthesis; metabolism of natural products; and detoxification of foreign substances (xenobiotics). *See e.g.*, Schuler, *Crit. Rev. Plant Sci.* 15:235-284 (1996). In a typical P-450  
15 catalyzed reaction, one atom of molecular oxygen (O<sub>2</sub>) is incorporated into the substrate, and the other atom is reduced to water by NADPH. For most eucaryotic P-450s, NADPH:cytochrome P-450 reductase, a membrane-bound flavoprotein, transfers the necessary two electrons from NADPH to the P-450 (Bolwell et al, *Phytochemistry* 37: 1491-1506 (1994)).

20 Frear et al. (*Phytochemistry* 8:2157-2169 (1969)) demonstrated the metabolism of monuron by a mixed-function oxidase located in a microsomal fraction of cotton seedlings. Further evidence has accumulated supporting the involvement of P-450s in the metabolism and detoxification of numerous herbicides representing several distinct classes of compounds (reviewed in  
25 Bolwell et al., 1994; Schuler, 1996). Differential herbicide metabolizing P-450 activities are believed to represent one of the mechanisms that enables certain crop species to be more tolerant of a particular herbicide than other crop or weedy species.

### Summary of the Invention

A first aspect of the present invention is an isolated DNA molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, or SEQ ID NO:17; or DNA sequences which encode an enzyme of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, or SEQ ID NO:18; or DNA sequences which have at least about 90% sequence identity to the above DNA and which encode a cytochrome P450 enzyme; and DNA sequences which differ from the above DNA due to the degeneracy of the genetic code.

A further aspect of the present invention is a cytochrome p450 enzyme having an amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, or SEQ ID NO:18.

A further aspect of the present invention is an isolated DNA molecule comprising SEQ ID NO:1; DNA sequences which encode an enzyme of SEQ ID NO:2; DNA sequences which have at least about 90% sequence identity to the above DNA and which encode a cytochrome P450 enzyme; and DNA sequences which differ from the above DNA due to the degeneracy of the genetic code.

A further aspect of the present invention is a cytochrome p450 peptide of SEQ ID NO:2.

A further aspect of the present invention is a DNA construct comprising a promoter operable in a plant cell and a DNA segment encoding a peptide of SEQ ID NO:2 downstream from and operatively associated with the promoter.

A further aspect of the present invention is a method of making a transgenic plant cell having an increased ability to metabolize phenylurea compounds compared to an untransformed plant cell. The plant cell is transformed with an exogenous DNA construct comprising a promoter operable in a plant cell and a DNA sequence encoding a peptide of SEQ ID NO:2.

Transformed plants, seed and progeny of such plants are also aspects of the

present invention.

A further aspect of the present invention is a transgenic plant having an increased ability to metabolize phenylurea compounds. Such transgenic plants contain exogenous DNA encoding a peptide of SEQ ID NO:2.

5

### **Brief Description of the Drawings**

**Figure 1** depicts dithionite-reduced carbon monoxide difference spectra, where the solid line represents microsomes isolated from yeast transformed with CYP71A10, and the dotted line shows the difference spectra from yeast  
10 transformed with control vector V-60. Microsomal protein concentration was 1 mg/ml.

**Figure 2** shows thin-layer chromatograms of [<sup>14</sup>C]-radiolabeled fluometuron, linuron, chlortoluron, and diuron and their respective metabolites after incubation of the radiolabeled herbicides with yeast microsomes containing  
15 the CYP71A10 protein. Initial substrate concentrations for fluometuron, linuron, chlortoluron and diuron were 5.2, 6.5, 4.0, and 3.7 μM, respectively. P = parent compound; M = metabolite.

**Figure 3** shows the chemical structures of fluometuron, linuron, chlortoluron and diuron, and their previously characterized metabolites. The  
20 linuron and chlortoluron metabolites are designated major or minor depending on their predicted relative abundance in assays using yeast microsomes containing the soybean CYP71A10 protein.

**Figure 4** shows thin-layer chromatograms using [<sup>14</sup>C]-radiolabeled linuron in various control reactions. The complete reaction mixture (COMPLETE)  
25 contained 3.2 μM linuron, 0.75 mM NADPH and 2.5 mg/ml microsomal protein isolated from CYP71A10-transformed yeast in 50 mM phosphate buffer (pH 7.1). Other reactions varied from COMPLETE by the addition of carbon monoxide (+CO), the omission of NADPH (NO NADPH), or the use of yeast microsomes isolated from cells expressing the control vector (V-60). P = parent  
30 compound; M = metabolite.

**Figure 5A** shows tobacco line 25/2 plants (transformed with soybean CYP71A10) grown on media containing no herbicide.

**Figure 5B** shows control tobacco plants (transformed with vector pBI121) grown on media containing 0.5  $\mu$ M linuron.

Figure 5C shows tobacco line 25/2 (transformed with soybean CYP71A10) individuals grown on media containing 0.5  $\mu$ M linuron.

Figure 5D shows tobacco line 25/2 (transformed with soybean CYP71A10) individuals grown on media containing 2.5  $\mu$ M linuron.

Figure 5E shows control tobacco plants (transformed with vector pBI121) grown on media containing 1.0  $\mu$ M chlortoluron.

Figure 5F shows tobacco line 25/2 (transformed with soybean CYP71A10) individuals grown on media containing 1.0  $\mu$ M chlortoluron.

15 Detailed Description of the Invention

### 1. Overview of the present research:

The present inventors utilized a strategy based on the random isolation and screening of soybean cDNAs encoding cytochrome P-450 (P-450) isozymes to identify P-450 isozymes involved in herbicide metabolism. Eight full-length and one near full-length P-450 cDNAs representing eight distinct P-450 families were isolated using polymerase chain reaction (PCR)-based technologies (SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15 and 17). Five of these soybean P-450 cDNAs were successfully overexpressed in yeast, and microsomal fractions generated from these strains were tested for their potential to mediate the metabolism of ten herbicides and one insecticide. *In vitro* enzyme assays showed that the gene product of one heterologously expressed P-450 cDNA (CYP71A10) (SEQ ID NO:1) specifically mediated the metabolism of phenylurea herbicides, converting four herbicides of this class (fluometuron, linuron, chlortoluron, and diuron) into more polar metabolites. Analyses of the metabolites indicate that the CYP71A10 encoded enzyme functions primarily as an N-demethylase with regard to

fluometuron, linuron and diuron, and as a ring-methyl hydroxylase when chlortoluron is the substrate. *In vivo* assays using excised leaves demonstrated that all four herbicides were more readily metabolized in CYP71A10-transformed tobacco in comparison to control plants.

5           Shiota et al. reported that fused constructs derived from the rat CYP1A1 and yeast NADPH-cytochrome P-450 oxidoreductase cDNAs conferred chlortoluron resistance in tobacco by enhancing herbicide metabolism (Shiota et al., *Plant Physiol.* 106:17-23 (1994)). In another study, a chloroplast-targeted, bacterial CYP105A1 expressed in tobacco catalyzed the toxification of R7402, a  
10           sulfonylurea pro-herbicide (O'Keefe et al., *Plant Physiol.* 105:473-482 (1994)). The cloning and heterologous expression of an endogenous plant P-450 gene that is potentially involved in herbicide metabolism was reported by Pierrel et al.,  
15           *Eur. J. Biochem.* 224:835-844 (1994), where a trans-cinnamic acid hydroxylase cDNA (CYP73A1) isolated from artichoke and expressed in yeast catalyzed the  
20           ring-methyl hydroxylation of chlortoluron. *In vivo* experiments with artichoke tubers, however, demonstrated that the ring-methyl hydroxy metabolite represented only a minor portion of the metabolites produced and that the major metabolite was demethylated chlortoluron (Pierrel et al., 1994). This together with the observation that the turnover number of the heterologously expressed  
25           enzyme was very low (0.014/ min), suggested that CYP73A1 plays a minimal role in chlortoluron metabolism *in vivo*. US Patent No. 5,349,127 to Dean et al. discloses the use of DNA encoding certain P-450 enzymes, isolated from *Streptomyces griseolus*, to produce transformed plants with increased metabolism of certain compounds. (All US patents referred to herein are intended to be  
30           incorporated herein in their entirety.)

          Although the role of P-450 enzymes in catalyzing the metabolism of a variety of herbicides has been documented, little progress has been made in the identification of the endogenous plant P-450s that are responsible for degrading these compounds. Protein purification of specific isozymes involved in the  
30           metabolism of a specific herbicide has been hindered by the instability of the

enzymes, their low concentrations in most plant tissues, and difficulties in the reconstitution of active complexes from solubilized components. Furthermore, any given plant tissue may possess dozens, if not hundreds, of unique P-450 isozymes, complicating the purification to homogeneity of a particular isozyme.

- 5 Because plants have only been exposed to phenylurea herbicides during the past few decades, it is unlikely that enzymes have evolved solely for the purposed of metabolizing this class of xenobiotics.

## 2. Use of CYP71A10 to produce phenylurea-resistant plants:

- 10 The present invention provides materials and methods useful in producing transgenic plant cells and plants with increased resistance to phenylurea herbicides. Increased herbicide resistance, as used herein, refers to the ability of a plant to withstand levels of an herbicide that have a negative impact on wild-type (untransformed) plants of the same species and/or variety. Resistance, as  
15 used herein, does not necessarily mean that the resistant plant is completely unaffected by exposure to the herbicide; rather, resistant plants suffer less extensive or less severe damage than comparable wild-type plants. Methods of assessing the extent and/or severity of herbicide impact will vary depending on the particular plant and the particular herbicide being tested; such assessment  
20 methods will be apparent to those skilled in the art. The negative effects of a herbicide may be evidenced by the complete arrest of plant growth, or by an inhibition in the rate or amount of growth. Additionally, methods of the present invention may be used to decrease herbicide residues in plants, even where the amounts of herbicides present in the plant do not cause an appreciable negative  
25 effect on the plant as a whole.

- Increased resistance to a herbicide can be due to an increased ability to metabolize a herbicide to less harmful metabolites. Accordingly, plants of the present invention which exhibit increased resistance to a herbicide may also be described as having an increased ability to metabolize the starting herbicidal  
30 compound, where the metabolites are less harmful to the plant than the starting



compound.

In the examples provided herein, yeast microsomes and transgenic tobacco plants expressing the CYP71A10 peptide (SEQ ID NO:2) and exposed to various phenylurea herbicides produced the same degradation products that have  
5 previously been observed when these same compounds have been incubated with metabolically active plant microsomes. These results indicate that the CYP71A10 peptide plays a role in the effective metabolism of phenylurea herbicides.

The present examples demonstrate that the overexpression of a  
10 CYP71A10 peptide of SEQ ID NO:2 in tobacco enhanced the plant's capacity to metabolize all four phenylurea herbicides tested, and that appreciable levels of tolerance were conferred to linuron and chlortoluron. Fluometuron was the most actively metabolized compound in both the yeast and transgenic plant systems, yet the enhancement in tolerance to this herbicide at the whole plant level was not  
15 as great as for linuron and chlortoluron. While not wishing to be held to a single theory, the present inventors surmise that the lack of correlation between the rate of herbicide metabolism and herbicide tolerance may be explained by the differential toxicities of the various phenylurea derivatives produced in the CYP71A10-transformed tobacco. Consistent with this hypothesis are the  
20 previous observations that N-demethyl derivatives of fluometuron, diuron and chlortoluron are only moderately less toxic than their parent compounds (Rubin and Eshel, *Weed Sci.* 19:592-594 (1971); Dalton et al., *Weeds* 14:31-33 (1966); Ryan and Owen, *Proc. Brit. Crop Prot. Conf. Weeds* 1:317-324 (1982)). In contrast, linuron is a 10-fold greater inhibitor of the Hill-reaction than N-  
25 demethyl linuron (Suzuki and Casida, *J. Agric. Food Chem.* 29:1027-1033 (1981)), and the hydroxylated and the didemethylated derivatives of chlortoluron are considered to be nonherbicidal (Ryan and Owen, 1982).

The present inventors found that the relative rates of herbicide metabolism in leaves of CYP71A10-transformed tobacco and in yeast microsomes assayed *in*  
30 *vitro* were similar (see Tables 4 and 5). With the exception of the transgenic

plant leaves showing a somewhat greater metabolic activity against chlortoluron than was apparent in the yeast microsomal assays, both systems followed the general order of metabolism of fluometuron  $\geq$  linuron  $>$  chlortoluron  $>$  diuron. These results indicate that expression of a test plant P-450 in yeast and  
5 quantification of the metabolism of a test compound using yeast microsomes, is a suitable system for screening plant P-450s for their metabolic function, and for their potential usefulness in the production of transgenic plants with altered metabolism of chemical compounds such as herbicides and insecticides.

The present inventors have shown that the random isolation of P-450  
10 cDNAs with subsequent heterologous expression in yeast is an effective strategy to characterize cDNAs whose product is capable of affecting the metabolism of a test compound. This approach is useful in characterizing the substrates (both natural and artificial) affected by a P-450, in determining the function of P-450 genes whose catalytic activities remain unclear, and in screening P-450s for the  
15 ability to increase or decrease the metabolism of a test compound. A particularly useful aspect of this method is the ability to screen isolated P-450s for their effects on the metabolism by plants of herbicides, insecticides, or other chemical compounds. Increased metabolism may result in enhanced resistance to the effects of a compound (where the metabolites are less harmful than the  
20 starting compound), or in increased sensitivity to the effects of a compound (where one or more metabolites are more toxic than the starting compound; *see* O'Keefe et al., 1994).

### 3. DNA Constructs:

25 Those familiar with recombinant DNA methods available in the art will recognize that one can employ a cDNA molecule (or a chromosomal gene or genomic sequence) encoding a P-450 peptide, joined in the sense orientation with appropriate operably linked regulatory sequences, to construct transgenic cells and plants. (Those of skill in the art will also recognize that appropriate  
30 regulatory sequences for expression of genes in the sense orientation include any

one of the known eukaryotic translation start sequences, in addition to the promoter and polyadenylation/transcription termination sequences described herein). Appropriate selection of the encoded P-450 peptide will provide transformed plants characterized by altered (enhanced or retarded) metabolism of phenylurea compounds.

DNA constructs, or "transcription cassettes," of the present invention include, 5' to 3' in the direction of transcription, a promoter as discussed herein, a DNA sequence as discussed herein operatively associated with the promoter, and, optionally, a termination sequence including stop signal for RNA polymerase and a polyadenylation signal for polyadenylase. All of these regulatory regions should be capable of operating in the cells of the tissue to be transformed. Any suitable termination signal may be employed in carrying out the present invention, examples thereof including, but not limited to, the nopaline synthase (nos) terminator, the octopine synthase (ocs) terminator, the CaMV terminator, or native termination signals derived from the same gene as the transcriptional initiation region or derived from a different gene. See, e.g., Rezian et al. (1988) *supra*, and Rodermel et al. (1988), *supra*.

The term "operatively associated," as used herein, refers to DNA sequences on a single DNA molecule which are associated so that the function of one is affected by the other. Thus, a promoter is operatively associated with a DNA when it is capable of affecting the transcription of that DNA (i.e., the DNA is under the transcriptional control of the promoter). The promoter is said to be "upstream" from the DNA, which is in turn said to be "downstream" from the promoter.

The transcription cassette may be provided in a DNA construct which also has at least one replication system. For convenience, it is common to have a replication system functional in *Escherichia coli*, such as ColE1, pSC101, pACYC184, or the like. In this manner, at each stage after each manipulation, the resulting construct may be cloned, sequenced, and the correctness of the manipulation determined. In addition, or in place of the *E. coli* replication

system, a broad host range replication system may be employed, such as the replication systems of the P-1 incompatibility plasmids, e.g., pRK290. In addition to the replication system, there will frequently be at least one marker present, which may be useful in one or more hosts, or different markers for individual hosts. That is, one marker may be employed for selection in a prokaryotic host, while another marker may be employed for selection in a eukaryotic host, particularly the plant host. The markers may be protection against a biocide, such as antibiotics, toxins, heavy metals, or the like; may provide complementation, by imparting prototrophy to an auxotrophic host; or may provide a visible phenotype through the production of a novel compound in the plant.

The various fragments comprising the various constructs, transcription cassettes, markers, and the like may be introduced consecutively by restriction enzyme cleavage of an appropriate replication system, and insertion of the particular construct or fragment into the available site. After ligation and cloning the DNA construct may be isolated for further manipulation. All of these techniques are amply exemplified in the literature as exemplified by J. Sambrook et al., Molecular Cloning, A Laboratory Manual (2d Ed. 1989)(Cold Spring Harbor Laboratory).

Vectors which may be used to transform plant tissue with nucleic acid constructs of the present invention include both *Agrobacterium* vectors and ballistic vectors, as well as vectors suitable for DNA-mediated transformation.

#### 4. Promoters:

The term 'promoter' refers to a region of a DNA sequence that incorporates the necessary signals for the efficient expression of a coding sequence. This may include sequences to which an RNA polymerase binds but is not limited to such sequences and may include regions to which other regulatory proteins bind together with regions involved in the control of protein translation and may include coding sequences.

Promoters employed in carrying out the present invention may be constitutively active promoters. Numerous constitutively active promoters which are operable in plants are available. A preferred example is the Cauliflower Mosaic Virus (CaMV) 35S promoter which is expressed constitutively in most plant tissues. Use of the CaMV promoter for expression of recombinant genes in tobacco roots has been well described (Lam et al., "Site-Specific Mutations Alter In Vitro Factor Binding and Change Promoter Expression Pattern in Transgenic Plants", *Proc. Nat. Acad. Sci. USA* 86, pp. 7890-94 (1989); Poulsen et al. "Dissection of 5' Upstream Sequences for Selective Expression of the *Nicotiana plumbaginifolia* rbcS-8B Gene", *Mol. Gen. Genet.* 214, pp. 16-23 (1988)). In the alternative, the promoter may be a tissue-specific promoter or a promoter that is expressed temporally or developmentally. See, e.g., US Patent No. 5,459,252 to Conkling et al.; Yamamoto et al., *The Plant Cell*, 3:371 (1991). In methods of transforming plants to alter the effects of herbicides or to decrease residual amounts of herbicides or pesticides in plants, selection of a suitable promoter will vary depending on the plant species, the specific chemical compound used as a herbicide or pesticide, and the time and method of applying the chemical compound to the plant or plant crop, as will be apparent to those skilled in the art.

#### 5. Selectable Markers:

The recombinant DNA molecules and vectors used to produce the transformed cells and plants of this invention may further comprise a dominant selectable marker gene. Suitable dominant selectable markers include, inter alia, antibiotic resistance genes encoding neomycin phosphotransferase (NPTII), hygromycin phosphotransferase (HPT), and chloramphenicol acetyltransferase (CAT). Another well-known dominant selectable marker suitable is a mutant dihydrofolate reductase gene that encodes methotrexate-resistant dihydrofolate reductase. DNA vectors containing suitable antibiotic resistance genes, and the corresponding antibiotics, are commercially available. Transformed cells are

selected out of the surrounding population of non-transformed cells by placing the mixed population of cells into a culture medium containing an appropriate concentration of the antibiotic (or other compound normally toxic to the untransformed cells) against which the chosen dominant selectable marker gene product confers resistance. Thus, only those cells that have been transformed will survive and multiply.

A further aspect of the present invention is use of the identified P-450 coding sequences as a selectable marker gene. A DNA construct comprising a sequence encoding a P-450 known to increase resistance to a compound (such as SEQ ID NO:2) is utilized to transform cells, in accordance with methods known in the art. Those cells that subsequently exhibit resistance to the compound are indicated as transformed. Such constructs may be used to verify the success of a transformation technique or to select transformed cells of interest.

15

#### 6. Sequence similarity and hybridization conditions:

Nucleic acid sequences employed in carrying out the present invention include those with sequence similarity to SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15 or 17, and encoding a protein having P-450 enzymatic activity. This definition is intended to encompass natural allelic variants and minor sequence variations in the nucleic acid sequence encoding a P-450 molecule, or minor sequence variations in the amino acid sequence of the encoded product. Thus, DNA sequences that hybridize to DNA of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15 or 17 and code for expression of a P-450 enzyme, particularly a plant P-450 enzyme, may also be employed in carrying out aspects of the present invention. The nomenclature for P-450 genes is based on amino acid sequence identity; methods of determining sequence similarity are well-known to those skilled in the art. Typically, sequences sharing >40% identity are placed in the same family, >55% identity defines members of the same subfamily, and sequences that

30

display > 97% identity are assumed to represent allelic variants. Conditions which permit other DNA sequences which code for expression of a protein having P-450 enzymatic activity to hybridize to DNA of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15 or 17, or to other DNA sequences encoding the protein given as  
5 SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16 or 18 can be determined in a routine manner. For example, hybridization of such sequences may be carried out under conditions of reduced stringency or even stringent conditions (e.g., conditions represented by a wash stringency of 0.3 M NaCl, 0.03 M sodium citrate, 0.1 % SDS at 60°C or even 70°C to DNA encoding the protein given as SEQ ID NO:2  
10 herein in a standard in situ hybridization assay. See J. Sambrook et al., Molecular Cloning, A Laboratory Manual (2d Ed. 1989)(Cold Spring Harbor Laboratory)). In general, such sequences will be at least 65% similar, 75% similar, 80% similar, 85% similar, 90% similar, 93% similar, 95% similar, or even 97% or 98% similar, or more, with the sequence given herein as SEQ ID  
15 NO:1, or DNA sequences encoding proteins of SEQ ID NO:2. (Determinations of sequence similarity are made with the two sequences aligned for maximum matching; gaps in either of the two sequences being matched are allowed in maximizing matching. Gap lengths of 10 or less are preferred, gap lengths of 5 or less are more preferred, and gap lengths of 2 or less still more preferred.)

20 As used herein, the term 'gene' refers to a DNA sequence that incorporates (1) upstream (5') regulatory signals including a promoter, (2) a coding region specifying the product, protein or RNA of the gene, (3) downstream (3') regions including transcription termination and polyadenylation signals and (4) associated sequences required for efficient and specific  
25 expression.

The DNA sequence of the present invention may consist essentially of a sequence provided herein (SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15 or 17), or equivalent nucleotide sequences representing alleles or polymorphic variants of these genes, or coding regions thereof.

30 Use of the phrase "substantial sequence similarity" in the present

specification and claims means that DNA, RNA or amino acid sequences which have slight and non-consequential sequence variations from the actual sequences disclosed and claimed herein are considered to be equivalent to the sequences of the present invention. In this regard, "slight and non-consequential sequence variations" mean that "similar" sequences (i.e., the sequences that have substantial sequence similarity with the DNA, RNA, or proteins disclosed and claimed herein) will be functionally equivalent to the sequences disclosed and claimed in the present invention. Functionally equivalent sequences will function in substantially the same manner to produce substantially the same compositions as the nucleic acid and amino acid compositions disclosed and claimed herein.

DNA sequences provided herein can be transformed into a variety of host cells. A variety of suitable host cells, having desirable growth and handling properties, are readily available in the art.

Use of the phrase "isolated" or "substantially pure" in the present specification and claims as a modifier of DNA, RNA, polypeptides or proteins means that the DNA, RNA, polypeptides or proteins so designated have been separated from their *in vivo* cellular environments through the efforts of human beings.

As used herein, a "native DNA sequence" or "natural DNA sequence" means a DNA sequence which can be isolated from non-transgenic cells or tissue. Native DNA sequences are those which have not been artificially altered, such as by site-directed mutagenesis. Once native DNA sequences are identified, DNA molecules having native DNA sequences may be chemically synthesized or produced using recombinant DNA procedures as are known in the art. As used herein, a native plant DNA sequence is that which can be isolated from non-transgenic plant cells or tissue.

#### 7. Transformed plants:

Methods of making recombinant plants of the present invention, in general, involve first providing a plant cell capable of regeneration (the plant cell



typically residing in a tissue capable of regeneration). The plant cell is then transformed with a DNA construct comprising a transcription cassette of the present invention (as described herein) and a recombinant plant is regenerated from the transformed plant cell. As explained below, the transforming step is carried out by techniques as are known in the art, including but not limited to bombarding the plant cell with microparticles carrying the transcription cassette, infecting the cell with an *Agrobacterium tumefaciens* containing a Ti plasmid carrying the transcription cassette, or any other technique suitable for the production of a transgenic plant.

10 Numerous *Agrobacterium* vector systems useful in carrying out the present invention are known. For example, U.S. Patent No. 4,459,355 discloses a method for transforming susceptible plants, including dicots, with an *Agrobacterium* strain containing the Ti plasmid. The transformation of woody plants with an *Agrobacterium* vector is disclosed in U.S. Patent No. 4,795,855. Further, U.S. Patent No. 4,940,838 to Schilperoort et al. discloses a binary *Agrobacterium* vector (i.e., one in which the *Agrobacterium* contains one plasmid having the vir region of a Ti plasmid but no T region, and a second plasmid having a T region but no vir region) useful in carrying out the present invention.

20 Microparticles carrying a DNA construct of the present invention, which microparticle is suitable for the ballistic transformation of a plant cell, are also useful for making transformed plants of the present invention. The microparticle is propelled into a plant cell to produce a transformed plant cell, and a plant is regenerated from the transformed plant cell. Any suitable ballistic cell transformation methodology and apparatus can be used in practicing the present invention. Exemplary apparatus and procedures are disclosed in Sanford and Wolf, U.S. Patent No. 4,945,050, and in Christou et al., U.S. Patent No. 5,015,580. When using ballistic transformation procedures, the transcription cassette may be incorporated into a plasmid capable of replicating in or  
25 integrating into the cell to be transformed. Examples of microparticles suitable  
30

-16-

for use in such systems include 1 to 5  $\mu\text{m}$  gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

Plant species may be transformed with the DNA construct of the present invention by the DNA-mediated transformation of plant cell protoplasts and subsequent regeneration of the plant from the transformed protoplasts in accordance with procedures well known in the art. Fusion of tobacco protoplasts with DNA-containing liposomes or via electroporation is known in the art. (Shillito et al., "Direct Gene Transfer to Protoplasts of Dicotyledonous and Monocotyledonous Plants by a Number of Methods, Including Electroporation", *Methods in Enzymology* 153, pp. 313-36 (1987)).

As used herein, transformation refers to the introduction of exogenous DNA into cells, so as to produce transgenic cells stably transformed with the exogenous DNA. Transformed plant cells are induced to regenerate intact plants through application of cell and tissue culture techniques that are well known in the art. The method of plant regeneration is chosen so as to be compatible with the method of transformation. The stable presence and the orientation of the exogenous DNA in transgenic plants can be verified by Mendelian inheritance of the DNA sequence, as revealed by standard methods of DNA analysis applied to progeny resulting from controlled crosses.

Plants of horticultural or agronomic utility, such as vegetable or other crops, can be transformed according to the present invention using techniques available in the art. A plant suitable for use in the present methods is *Nicotiana tabacum*, or tobacco. Any strain or variety of tobacco may be used. Additional plants (both monocots and dicots) which may be employed in practicing the present invention include, but are not limited to, potato (*Solanum tuberosum*), soybean (*Glycine max*), tomato (*Lycopersicon esculentum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium hirsutum*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus spp.*), cassava (*Manihot esculenta*), coffee (*Cofea spp.*), pineapple (*Ananas comosus*), citrus trees (*Citrus*

spp.), banana (*Musa* spp.), corn (*Zea mays*), oilseed rape (*Brassica napus*), wheat, oats, barley, rye and rice. Thus, an illustrative category of plants which may be used to practice aspects of the present invention are the dicots, and a more particular category of plants which may be used to practice the present invention are members of the family Solanaceae.

The methods of the present invention can further be practiced with turfgrass, including cool season turfgrasses and warm season turfgrasses. Examples of cool season turfgrasses are Bluegrasses (*Poa* L.), such as Kentucky Bluegrass (*Poa pratensis* L.), rough Bluegrass (*Poa trivialis* L.), Canada Bluegrass (*Poa compressa* L.), Annual Bluegrass (*Poa annua* L.), Upland Bluegrass (*Poa glaucantha* Gaudin), Wood Bluegrass (*Poa nemoralis* L.), and Bulbous Bluegrass (*Poa bulbosa* L.); the Bentgrasses and Redtop (*Agrostis* L.), such as Creeping Bentgrass (*Agrostis palustris* Huds.), Colonial Bentgrass (*Agrostis tenius* Sibth.), Velvet Bentgrass (*Agrostis canina* L.), South German Mixed Bentgrass (*Agrostis* L.), and Redtop (*Agrostis alba* L.); the Fescues (*Festuca* L.), such as Red Fescue (*Festuca rubra* L.), Chewings Fescue (*Festuca rubra* var. *commutata* Gaud.), Sheep Fescue (*Festuca ovina* L.), Hard Fescue (*Festuca ovina* var. *duriuscula* L. Koch), Hair Fescue (*Festuca capillata* Lam.), Tall Fescue (*Festuca arundinacea* Schreb.), Meadow Fescue (*Festuca elatior* L.); the Rye grasses (*Lolium* L.), such as Perennial Ryegrass (*Lolium perenne* L.), Italian Ryegrass (*Lolium multiflorum* Lam.); the Wheatgrasses (*Agropyron* Gaertn.), such as Fairway Wheatgrass (*Agropyron cristatum* L. Gaertn.), Western Wheatgrass (*Agropyron smithii* Rydb.). Examples of warm season turfgrasses are the Bermudagrasses (*Cynodon* L.C. Rich), the Zoysiagrasses (*Zoysia* Willd.), St. Augustinegrasses (*Stenotaphrum secundatum* (Walt.) Kuntze), Centipedegrass (*Eremochioa ophiuroides* (Munro.) Hack.), Carpetgrass (*Axonopus* Beauv.), Bahiagrass (*Paspalum notatum* Flugge.), Kikuyugrass (*Pennisetum clandestinum* Hochst. ex Chiov.), Buffalograss (*Buchloe dactyloides* (Nutt.) Engelm.), Blue Grama (*Bouteloua gracilis* (H.B.K.) Lag. ex Steud.), Sideoats Grama (*Bouteloua curtipendula* (Michx.) Torr.), and Dichondra

(*Dichondra* Forst.).

Any plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a vector of the present invention. The term "organogenesis," as used herein, means a process by which shoots and roots are developed sequentially from meristematic centers; the term "embryogenesis," as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, callus tissue, existing meristematic tissue (e.g., apical meristems, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem).

Plants of the present invention may take a variety of forms. The plants may be chimeras of transformed cells and non-transformed cells; the plants may be clonal transformants (e.g., all cells transformed to contain the transcription cassette); the plants may comprise grafts of transformed and untransformed tissues (e.g., a transformed root stock grafted to an untransformed scion in citrus species). The transformed plants may be propagated by a variety of means, such as by clonal propagation or classical breeding techniques. For example, first generation (or T1) transformed plants may be selfed to provide homozygous second generation (or T2) transformed plants, and the T2 plants further propagated through classical breeding techniques. A dominant selectable marker (such as nptII) can be associated with the transcription cassette to assist in breeding.

As used herein, a crop comprises a plurality of plants of the same genus or species, planted together in an agricultural field. By "agricultural field" is meant a common plot of soil or a greenhouse. Thus, the present invention provides a method of producing a crop of plants having altered metabolism of chemical compounds (such as a phenylurea herbicide), and thus having altered

resistance to the chemical compound, compared to a crop of non-transformed plants of the same genus or species, or variety.

Where a crop comprises a plurality of transgenic plants with increased resistance to phenylurea compounds according to the present invention, such compounds may be used as post-emergent herbicides to control undesirable plant species. Accordingly, a method of using phenylurea compounds as post-emergent herbicides according to the present invention comprises planting a plurality of transformed plant seed (or transformed plants) with enhanced resistance to a phenylurea herbicide, and applying that herbicide to the field after the germination and emergence of at least some of said transformed plant seed (or following the planting of transformed plants). Application of the phenylurea herbicide will selectively impact non-resistant plants.

#### 9. Microbial decontamination:

Microbial cells useful for degrading phenylurea compounds, which cells contain and express a heterologous DNA molecule encoding a P-450 enzyme that enhances the metabolism of the phenylurea compound in the microbial cell (*e.g.*, a peptide of SEQ ID NO:2), are a further aspect of the present invention. Suitable host microbial cells include soil microbes (*i.e.*, those which grow in the soil) transformed to express a P-450 enzyme that enhances the metabolism of one or more phenylurea compounds by the host cell. Suitable microbes include bacteria (such as *Agrobacterium*, *Bacillus*, *Streptomyces*, *Nocardia*, etc.), fungi (including yeasts), and algae. Microbes can be selected, by methods known in the art of soil microbiology, to correspond to those which are typically found in the substrate to be treated. Liquids which are contaminated with phenylurea compounds may be contacted to transformed microorganisms by passing the contaminated liquid through a bioreactor which contains the microorganism. Numerous suitable bioreactor designs are known in the art. A microbial host particularly suitable for bioreactors is yeast.

Combination treatments utilizing aspects of the present invention involve

the application of a phenylurea compound in a location such as an agricultural field (*e.g.*, as a herbicide), and subsequent application of a transformed microbe as described above in an amount effective to degrade residual applied herbicide. Application of the herbicide may be carried out in accordance with known techniques.

The examples which follow are set forth to illustrate the present invention, and are not to be construed as limiting thereof.

## EXAMPLE 1

10

### Materials and Methods

#### a. Substrates

Phenyl-U-[<sup>14</sup>C] fluometuron, phenyl-U-[<sup>14</sup>C] chlortoluron, phenyl-U-[<sup>14</sup>C] metolachlor, phenyl-U-[<sup>14</sup>C] prosulfuron, pyrimidinyl-2- diazinon, and phenyl-U-[<sup>14</sup>C] alachlor were provided by Novartis (Greensboro, North Carolina); phenyl-U-[<sup>14</sup>C] bentazon was donated by BASF (Research Triangle Park, North Carolina); phenyl-U-[<sup>14</sup>C] linuron, phenyl-U-[<sup>14</sup>C] diuron, and carbonyl-[<sup>14</sup>C] metribuzin were a gift from DuPont de Nemours (Wilmington, Delaware); carboxyl-[<sup>14</sup>C] imazaquin was provided by American Cyanamid (Princeton, New Jersey).

20

#### b. Isolation of P-450 cDNAs

Random amplification of partial cDNAs encoding P-450 enzymes was conducted essentially as described by Meijer et al., *Plant Mol. Biol.* 22:379-383 (1993), using a soybean (*Glycine max* cv Dare) leaf cDNA library as the template (Dewey et al., *Plant Cell* 6:1495-1507 (1994)). Briefly, degenerate inosine-containing primers were synthesized based on the highly conserved heme-binding region. The precise sequences of these primers are described in Meijer et al. (1993). An oligo-dT primer complementary to the poly(A) tail of the cDNA clones was used in conjunction with the degenerate primers in PCR amplification assays. Amplification products were cloned into the T-tailed pCRII plasmid

(Invitrogen, San Diego, CA) and DNA sequence analysis of the first 300-400 base pairs downstream of the conserved region was used to establish whether a given amplification product represented a true P-450 cDNA.

To recover full-length versions of the partial cDNAs, a primer (5'-  
5 TGTCTAACTCCTTCCTTTTC-3') (SEQ ID NO:19) complementary to the pYES2 vector (the vector into which the soybean cDNA library was cloned) and a downstream primer corresponding to a segment of the 3' untranslated region for each of the unique P-450 cDNAs were used in PCR reactions using the same soybean cDNA library as the template. PCR products were again cloned into the  
10 pCRII plasmid and the entire DNA sequence was determined for the largest cDNA amplified for each unique soybean P-450.

To isolate full-length versions of the respective P-450 ORFs without including any of the 5' untranslated region (which has been shown to potentially impede gene expression in yeast (Pompon, *Eur. J. Biochem.* 177:285-293  
15 (1988)), an additional PCR reaction was performed with two gene-specific primers. The forward primers contained a BamHI restriction site immediately followed by the ATG start codon, and the next 14-15 bases of the reading frame; the downstream primer was again specific for the 3' untranslated regions of the respective genes and included sequences specifying either EcoRI, KpnI, and SacI  
20 to facilitate subcloning of the P-450 cDNAs into the yeast expression vector, pYeDP60 (V-60; Urban et al., *Biochimie* 72:463-472 (1990)).

All PCR reactions, with the exception of the initial amplification of the partial P-450 cDNAs (see Meijer et al. (1993)), contained 0.2 ng/ $\mu$ l template, 2  $\mu$ M of each primer, 200  $\mu$ M of each dNTP, and 1.5 mM  $MgCl_2$  in a final  
25 reaction volume of 50  $\mu$ l. Amplification was initiated by the addition of 1.5 U EXPAND<sup>TM</sup> High Fidelity enzyme mix using conditions described by the manufacturer (Boeringer Mannheim). DNA sequence was determined by the chain termination method (Sanger et al., *Proc. Natl. Acad. Sci. USA* 74:5463-5467 (1977)) using fluorescent dyes (Applied Biosystems, Foster City, CA).  
30 DNA and predicted amino acid sequences were analyzed using the BLAST

algorithm and the GAP program (University of Wisconsin, Madison, Genetics Computing Group software package).

c. P-450 cDNA Expression in Yeast

5 Yeast transformation was performed as described by Geitz et al., *Nucleic Acids Research* 20:1425 (1992). Media composition, culturing conditions, galactose induction, and microsomal preparations were conducted according to Pompon et al., *Methods Enzymol.* 272:51-64 (1995), using a culture volume of 250 ml. Microsomal protein was quantified spectrophotometrically using the  
10 method of Waddell, *J. Lab. Clin. Med.* 48:311-314 (1956), using bovine albumin as a standard. Dithionite-reduced, carbon monoxide difference spectra was obtained as previously outlined (Estabrook and Werringloer, *Methods Enzymol.* 52:212-220 (1978)) using a Shimadzu Recording Spectrophotometer UV-240 (Shimadzu, Kyoto, Japan). P-450 protein concentrations of yeast microsomes  
15 were calculated using a millimolar extinction coefficient of 91 (Omura and Sato, *J. Biol. Chem.*, 239:2370-2378 (1964)).

d. In vitro Herbicide Metabolism Assays

Yeast microsomes enriched for a discrete soybean P-450 isozyme were  
20 assayed for their capacity to metabolize the ten herbicides and one insecticide listed in Table 3. The reaction mixtures contained 10,000 DPM (100-200 ng) radiolabeled substrate, 0.75 mM NADPH, 2.5 mg/ml microsomal protein. Total reaction volumes were adjusted to 150  $\mu$ l with 50 mM phosphate buffer (pH 7.1).

The mixtures were incubated under light for 45 minutes at 27°C, arrested with  
25 50  $\mu$ l acetone and centrifuged at 14 000xg for 2 minutes. Fifty microliters of the supernatants containing radiolabeled alachlor, metolachlor, metribuzin, prosulfuron, chlortoluron, diuron, fluometuron, linuron, or diazinon were spotted onto 250 micron Whatman K6F silica plates. Radiolabeled bentazon and imazaquin-containing samples were spotted onto 200 micron Whatman LKC18F  
30 silica gel reversed-phase plates. All plates were developed in a benzene/acetone



2:1 (v/v) solvent system with the exception of prosulfuron, developed in toluene/acetone/acetic acid, 75:20:5 (v/v/v), and bentazon and imazaquin, developed in methanol/75 mM sodium acetate 40:60 (v/v). The developed plates were scanned with a Bioscan System 400 imaging scanner (Bioscan, Washington, DC), and the production of metabolites was determined based on the chromatographic profiles. For microsomes containing the expressed CYP71A10 enzyme, control experiments were also conducted to measure the NADPH-dependency, and the inhibitory effects of CO. CO treatment of the sample was achieved by gentle bubbling of the gas through the reaction mixture for 2 minutes immediately before the assay was initiated by the addition of NADPH.

#### e. Enzyme Kinetics

Substrate conversion was quantified by a combination of TLC analysis and scintillation spectrometry. The location of the metabolic products on the TLC plates was identified using an imaging scanner, the bands were scraped and analyzed by scintillation spectrometry. The amount of metabolite produced was calculated based on specific activity and scintillation counts. Each assay was repeated at least twice.  $K_m$  and  $V_{max}$  values were estimated using nonlinear regression analysis.

#### f. Mass Spectral Analysis

The reaction components used in the *in vitro* fluometuron and linuron metabolism assays were scaled up 50-fold, and the reactions were allowed to proceed for 3 hours. The substrates and the metabolites were extracted 3 times with 20 ml ethyl acetate. The extracts were combined, evaporated to dryness, and the resulting pellet was resuspended in 1 ml acetone. The samples were purified twice using preparative TLC and imaging scanning as described above. Finally, the respective bands were scraped, the compounds were eluted with acetone and flash evaporated.

Fractions of interest were analyzed by liquid chromatography/mass

spectrometry (LC/MS). Mass spectral measurements were made with a Finnigan TSQ 7000 triple quadrupole mass spectrometer (QQQ) equipped with an Atmospheric Pressure Ionization (API) interface fitted with a pneumatically assisted electrospray head (Finnigan MAT, Bremen, Germany). The spray  
5 nozzle was operated at 5 kV in the positive ion mode and 4 kV in the negative ion mode. For sample introduction, the TSQ 7000 was equipped with a HPLC solvent delivery system (Perkin-Elmer 410 LC pump), a UV detector (Perkin-Elmer), a stream splitter set at 6:1 with the majority of the effluent flowing to a radioisotope flow monitor (IN/US  $\beta$ -RAM) and the other stream attached to the  
10 API interface. Samples were chromatographed on a reverse phase HPLC column (Inertsil 5 ODS2, 150 x 2 mm i.d.). The column was eluted at 0.4 ml/min with 95:5 of 0.1% trifluoroacetic acid in water and 0.1% trifluoroacetic acid in methanol, respectively. Collision induced dissociation experiments (MS/MS) were conducted using argon gas with collision energy in the range of 17.5-30 eV  
15 at cell pressures of approximately 0.28 Pa. Signals were captured using a Finnigan 7000 data system.

#### g. NMR Analysis

Proton NMR measurements were made on a Bruker AMX-400 NMR  
20 spectrometer equipped with either a QNP or inverse probe set at 400.13 MHz. Spectra were acquired at ambient temperature in acetonitrile- $d_3$ . Chemical shifts were expressed as parts per million, relative to the resonance of residual acetonitrile protons at 1.93 ppm ( $\delta$ ).

#### 25 h. Tobacco Transformation

A plant expression vector capable of mediating the constitutive expression of CYP71A10 was produced. The GUS open reading frame of the binary expression vector pBI121 (Clontech, Palo Alto, CA) was excised and replaced with the full length CYP71A10 reading frame. This placed the soybean gene  
30 under the transcriptional control of the strong constitutive CaMV 35S promoter.

The resulting construct was used to transform *Agrobacterium tumefaciens* strain LBA 4404 (Holsters et al., *Mol. Gen. Genetics*, 163:181-187 (1988)). Excised leaf discs of *Nicotiana tabacum* cv SR1 were transformed using the *Agrobacterium*, and kanamycin-resistant plants were selected as described by  
5 Horsch et al. *Science*, 227:1229-1231 (1985). Primary transformants were potted in a standard soil mixture, transferred to a greenhouse and their seed harvested upon maturation.

i. In vivo Herbicide Metabolism Assays

10 Seeds from primary transgenic tobacco plants transformed with CYP71A10 and control plants transformed with the pBI121 vector were grown in Petri dishes containing MS salts and 100 µg/ml kanamycin. At five weeks post-seeding, kanamycin-resistant plantlets were transplanted into pots containing soil and grown an additional two weeks. Single leaves of approximately 10 cm<sup>2</sup> in  
15 size were excised and their petioles inserted into 100 µl of H<sub>2</sub>O containing radiolabeled herbicide. The leaves were placed in a growth chamber maintaining a temperature of 27°C and incubated until the entire volume of the herbicide solution was drawn up by the transpirational stream of the leaves (about 3 hrs). The leaves were subsequently transferred into an Eppendorf tube containing  
20 distilled water and further incubated for a total of 14 hours.

[<sup>14</sup>C]-labeled herbicide was extracted from the leaves by grinding for 5 minutes in 250 µl methanol with a plastic pellet pestle driven by an electric drill. After centrifugation for 3 minutes at 14,000 g, 75µl of the supernatant was spotted on a Whatman K6F silica plate and developed in a solvent system  
25 containing chloroform/ethanol/acetic acid 135:10:15 (v/v/v). The separated herbicide derivatives were visualized using an imaging scanner. Substrate conversion was quantified based on the amount of herbicide absorbed, and the ratios of the parent compound and the produced metabolites determined from the TLC profiles.

30

#### j. Herbicide Tolerance

T<sub>1</sub> generation seeds from CYP71A10-transformed tobacco and pBI121-transformed control plants were placed onto Petri dishes containing MS salts and linuron (using its commercial formulation LOROX 50 DF) at active ingredient concentrations ranging from 0.25 to 3.0  $\mu$ M. Chlortoluron was added at 0, 1.0, 5.0 and 10.0  $\mu$ M concentrations using a 99.5% pure analytical standard. The Petri dishes were incubated in a growth chamber maintaining a constant temperature of 27°C and a 16/8 hour light/dark cycle. The phytotoxic effects of the treatments were determined visually by comparison to control plants and plants grown in the absence of the herbicide. All treatments were repeated at least twice.

#### EXAMPLE 2

##### Isolation of P-450 cDNAs

To isolate cDNAs encoding P-450s from soybean, the PCR strategy described by Meijer et al. (1993) was adapted, using a soybean leaf cDNA library as the template. Degenerate, inosine-containing PCR primers were constructed corresponding to the first nine codons encoding the conserved sequence FLPGGxGxRxG (x = any amino acid) (SEQ ID NO:20), which represents an extension of the highly conserved FxxGxxxG motif (Bozak et al., *Proc. Natl. Acad. Sci. USA* 87:3904-3908 (1990)) (SEQ ID NO:21). Located near the C-terminal end of the protein, this motif defines the heme-binding region of the protein and may be regarded as a "signature" for P-450 proteins. A second nonspecific primer complementary to the poly(A) tail of the cDNA clones was used in conjunction with these degenerate primers in a PCR amplification assay. PCR amplification products were cloned into a plasmid vector and analyzed by DNA sequencing. Of 86 randomly selected individuals that were sequenced, 15 clones representing 10 unique cDNAs were identified that possessed the conserved cysteine and glycine residues of the signature

consensus (xCxG) (SEQ ID NO:22) immediately following the sequence defined by the degenerate PCR primers. Furthermore, homology searches of the major DNA and protein data bases revealed additional sequence identities to previously reported P-450 sequences for each of the ten unique soybean sequences (data not shown). Because this strategy only allows the recovery of sequence corresponding to the C-terminal portion of the proteins, additional PCR-based techniques were utilized to obtain cDNAs possessing the entire reading frames for each clone. Full length cDNAs were isolated for eight of the 10 individual clones and a near full length cDNA was isolated for an additional clone.

10       The eight full length and one near full length soybean P-450 cDNAs isolated are described in Table 1. The nomenclature for P-450 genes is based on amino acid sequence identity. Typically, sequences sharing >40% identity are placed in the same family, >55% identity defines members of the same subfamily, and sequences that display >97% identity are assumed to represent  
15       allelic variants, although exceptions to these designations have been noted (Nelson et al., *Pharmacogenetics*, 6:1-41 (1996)). According to this system of nomenclature, all of the nine soybean cDNAs were able to be placed within existing P-450 gene families; however, three of the sequences (CYP82C1, CYP83D1 and CYP93C1) defined new subfamilies. Although an increasing  
20       number of P-450 gene products have been assigned specific enzymatic functions (reviewed in Schuler, 1996), none of the soybean cDNAs listed in Table 1 could be placed into families for which an *in vivo* function had been determined for any of its members.

      In addition to the conserved heme-binding domain described previously,  
25       all of the predicted soybean polypeptides possess slight variations of the conserved sequence PEEFxPERF (SEQ ID NO:23) located approximately 30 amino acids forward of the heme-binding motif (Hallahan et al., *Biochem. Soc. Trans.* 21:1068-1073 (1993)). Also characteristic of microsomal P-450s is the presence of an N-terminal noncleavable signal sequence that serves as the  
30       membrane anchor. Immediately following this signal-anchor segment in most

microsomal P-450s is a proline-rich region that is believed to form a hinge between the catalytic cytoplasmic domain and the hydrophobic membrane anchor (Halkier, *Phytochemistry* 43:1-21 (1996)). All of the present clones (except CYP97B2) encode proteins possessing predicted signal sequences; all individuals (except CYP97B2 and CYP82C1) contain readily identifiable proline-rich domains following the signal sequence (Table 1). It is the identification of both of these N-terminal motifs in the CYP83D1 encoded protein (but no Met codon) that indicates that this clone is nearly full length. Interestingly, instead of possessing a predicted signal sequence and proline-rich region, the N-terminus of the polypeptide encoded by clone CYP97B2 contains a motif characteristic of a chloroplast transit peptide (data not shown).

**Table 1**  
**Soybean P-450s Isolated Using Degenerate PCR Primers**

Name	GenBank Accession #	Length (amino acids)	Closest Match	Identity* %	Membrane Anchor	Proline-rich Region
CYP71A10 (SEQ ID NO:1)	AF022157	513	CYP71A1	51.7	+	+
CYP71D10 (SEQ ID NO:3)	AF022459	510	CYP71D9	50.9	+	+
CYP77A3 (SEQ ID NO:5)	AF022464	513	CYP77A1	69.8	+	+
CYP78A3 (SEQ ID NO:7)	AF022463	523	CYP78A2	53.1	+	+
CYP82C1 (SEQ ID NO:9)	AF022461	532	CYP82A3	51.1	+	-
CYP83D1** (SEQ ID NO:11)	AF022460	516	CYP71A1**	45.7	+	+
CYP93C1 (SEQ ID NO:13)	AF022462	521	CYP93B1	44.5	+	+
CYP97B2 (SEQ ID NO:15)	AF022457	576	CYP97B1	80.8	-	-
CYP98A2 (SEQ ID NO:17)	AF022458	509	CYP98A1	69.7	+	+

\*Percent identity between the predicted amino acids sequences of the given soybean P-450 cDNA and the closest match identified from a BLAST search against the major gene and protein databases.

\*\* Although this sequence shows a best match to CYP71A1, it matches poorly to some sequences of the CYP71B subfamily. As a result, the tree cluster program places it into the CYP83 family.

## EXAMPLE 3

## Expression of Soybean P-450 cDNAs in Yeast

Because superfluous 5' untranslated sequences from foreign genes have  
5 been shown to be capable of impeding gene expression in yeast (Pompon, 1988),  
an additional PCR reaction was performed on each clone that enabled the  
cloning of full length P-450 open reading frames (ORFs) into the yeast  
expression vector pYeDP60 (V-60) without including any of the endogenous 5'  
nontranslated flanking sequence (see Methods). For the near full length clone  
10 CYP83D1, the 5' primer was also designed to generate an "artificial" Met start  
codon and a Val second codon at the 5' end of the ORF. Expression in yeast of  
genes cloned into the V-60 vector is mediated by the strong, galactose-inducible  
GAL10-CYC1 promoter (Pompon et al., 1995).

Previous studies have revealed that the heterologous expression of P-450  
15 cDNAs in yeast can be greatly enhanced in strains that have been engineered to  
overexpress endogenous NADPH-dependent cytochrome P-450 reductase  
(Pompon et al., 1995). In strain W(R), this was accomplished by exchanging the  
relatively weak endogenous cytochrome P-450 reductase promoter with the same  
GAL10-CYC1 promoter used in vector V-60 (Truan et al., *Gene* 125:49-55  
20 (1993)). To maximize the heterologous expression of the soybean P-450 cDNAs  
in yeast, each of the constructs cloned into the V-60 vector was transformed into  
strain W(R) and microsomes were isolated from cultures that had been induced  
by galactose.

Reduced-CO difference spectroscopy provides a method to measure the  
25 effectiveness of expression of heterologous P-450s in yeast. Microsomal  
preparations corresponding to five of the soybean constructs (CYP71A10,  
CYP71D10, CYP77A3, CYP83D1 and CYP98A2) showed characteristic P-450  
CO difference spectra with Soret peaks at 450 nm; the profile corresponding to  
CYP71A10 is shown in Figure 1. No such peaks were observed for the  
30 remaining four clones. The specific P-450 content of the five positive

microsomal preparations varied significantly, ranging from 11 pmol P-450/mg protein for construct CYP83D1 to 252 pmol P-450/mg for clone CYP77A3 as shown in Table 2.

5

**Table 2**  
**P-450 Content of Microsomes Isolated from Yeast Overexpressing Various Soybean CYPs**

Clone	CYP content (pmol mg <sup>-1</sup> protein)
CYP71A10	44
CYP71D10	15
CYP77A3	252
CYP83D1	11
CYP98A2	13

10

#### EXAMPLE 4

##### *In vitro* Herbicide Assays

To determine whether any of the present soybean P-450 proteins synthesized in yeast displayed significant herbicide metabolic activity, 15 microsomal preparations possessing each of the five soybean P-450s that were effectively expressed in yeast (as judged by their reduced CO difference spectra, see above) were incubated individually with NADPH and radioisotopes of the compounds listed in Table 3. These substrates represent six different classes of herbicides and one organophosphate insecticide (diazinon). Upon termination of 20 the reaction, each sample was analyzed by thin layer chromatography (TLC) to reveal potential metabolic breakdown products.

The P-450 proteins expressed from clones CYP71D10, CYP77A3, CYP83D1, and CYP98A2 displayed no apparent *in vitro* metabolic activity against any of the 11 compounds tested (data not shown). In contrast, the P-450 25 enzyme produced from construct CYP71A10 demonstrated considerable activity



against the phenylurea class of herbicides, but no activity against the remaining compounds. As shown in Figure 2, fluometuron and diuron were converted to a single metabolite; linuron and chlortoluron were transformed into two (a major and a minor) metabolites. Figure 3 shows the chemical structures of the four phenylurea herbicides tested in this study, and the derivatives that have previously been characterized as the first metabolites produced during the detoxification of the respective herbicides in plants known to metabolize these compounds (Voss and Geissbühler, *Proc. Brit. Weed Contr. Conf.* 8:266-268 (1966); Suzuki and Casida, *J. Agric. Food Chem.* 29:1027 (1981); Ryan et al., *Pestic. Biochem. Physiol.* 16:213-221 (1981)).

To further confirm that the herbicide metabolism measured from microsomes of yeast expressing CYP71A10 was attributable to a P-450 activity, additional assays utilizing linuron as the substrate were conducted. As shown in Figure 4, linuron metabolizing activity is reduced approximately 37% in the presence of CO, and no metabolites are observed when NADPH is omitted from the reaction. Activity is also completely abolished upon addition of tetracycline, a potent P-450 inhibitor (data not shown). Furthermore, no activity is detected when microsomal preparations are used from yeast cells expressing only the V-60 control plasmid. These results verify that the observed herbicide metabolizing activity is derived from the soybean CYP71A10 cDNA.

The kinetic properties and catalytic activities of the soybean CYP71A10 protein enzyme differed significantly among the four phenylurea-type herbicide substrates. As shown in Table 4, turnover rates for fluometuron and linuron were considerably greater than those observed for chlortoluron and diuron. The observed reduced activity for the later two substrates is apparently not the result of decreased binding affinities since the apparent  $K_m$ s for chlortoluron and diuron are lower than those measured for fluometuron and linuron.

**Table 3**

**30 Compounds Used in Metabolism Assays**

Common Name	Chemical Family
Alachlor	Acetanilide
Metolachlor	Acetanilide
Bentazon	Benzothiadiazole
Imazaquin	Imidazolinone
Chlortoluron	Phenylurea
Diuron	Phenylurea
Fluometuron	Phenylurea
Linuron	Phenylurea
Prosulfuron	Sulfonylurea
Metribuzin	<i>as</i> -Triazine
Diazinon	Organophosphate

**Table 4**  
**In Vitro Kinetic Parameters of the CYP71A10 Enzyme**  
**for Four Phenylurea Substrates**

Substrate	$K_{m, app}$	$V_{max}$	Turnover
	( $\mu M$ )	( $pmol\ min^{-1}\ mg^{-1}$ protein)	( $min^{-1}$ )
Fluometuron	14.9 (1.0)*	303.6 (10.8)	6.8 (0.24)
Linuron	9.8 (2.1)	125.6 (12.0)	2.8 (0.27)
Chlortoluron	1.0 (0.2)	29.4 (2.2)	0.7 (0.05)
Diuron	1.5 (0.3)	16.8 (1.6)	0.4 (0.04)

- 5    \* Values in parentheses represent standard error.  
      - Assays were repeated three times for linuron and twice for all other substrates.  
      - Concentration ranges ( $\mu M$ ) used were 3.2-27.7 for fluometuron, 3.8-28.3 for linuron, 0.7-4.0 for chlortoluron, and 0.7-3.7 for diuron.

10

### EXAMPLE 5

#### Analysis of Metabolites

As shown in **Figure 2**, CYP71A10-mediated degradation of phenylurea herbicides resulted in the accumulation of either one or two metabolites, depending on the particular substrate used. To determine the structure of the metabolites, the single metabolite observed in the fluometuron assay and both the major and minor metabolites generated in the linuron assay were analyzed by liquid chromatography/mass spectroscopy (LC/MS) analysis (results not shown).

Analysis of the fluometuron metabolite by LC/MS in positive ion mode resulted in pseudomolecular ions at  $m/z$  219  $[(M+H)^+, C_9H_9F_3N_2O]$  and  $m/z$  241  $(M+Na)^+$  that corresponds to a sodium adduct. Daughter ion spectra of  $m/z$  219 produced a prominent  $m/z$  162 fragment ion due to formation of the protonated trifluoromethylaniline ( $C_7H_6F_3N+H$ ) $^+$ . Analysis of the fluometuron metabolite by proton NMR showed a singlet at  $\delta$ 2.71 which integrated for 3 protons (data not shown). The NMR spectra aromatic resonances were similar to aromatic resonances observed in the parent molecule. Spectra of the fluometuron metabolite were consistent for loss of a methyl group from the parent compound.

The major linuron metabolite analyzed by LC/MS in the negative ion mode showed a pseudomolecular ion at  $m/z$  233  $(M-H)^-$  and  $m/z$  235  $[(M+2)-H]^+$  consistent for a molecule containing two chlorine atoms. Daughter ion spectrum at  $m/z$  233 showed a prominent fragment ion at  $m/z$  160  $(C_6H_4Cl_2N-H)^+$ . The major linuron metabolite was 15 mass units less than parent compound which is consistent with loss of a methyl group. The position of methyl loss could not be determined based on mass spectral data alone.

The minor linuron metabolite analyzed by LC/MS gave a pseudomolecular ion at  $m/z$  217  $(M-H)^-$  and  $m/z$  219  $[(M+2)-H]^+$  which is consistent for a molecule containing two chlorine atoms. The daughter ion spectrum at  $m/z$  217 showed a prominent fragment ion at  $m/z$  160 which corresponds to formation of the dichloroaniline. The mass spectral data is consistent for the minor linuron metabolite representing N-demethoxy linuron.

These results suggest that the CYP71A10 enzyme expressed in yeast produces the same fluometuron and linuron metabolites as depicted in Figure 3, which shows the first metabolites produced during the detoxification of the respective herbicides in plants that are known to degrade these compounds. The metabolites of chlortoluron and diuron have not been analyzed directly, but the  $R_f$  values of the peaks observed during TLC separation are consistent with these species also representing the compounds shown in Figure 3 (ring-hydroxymethyl chlortoluron, N-demethyl chlortoluron and N-demethyl diuron). These results indicate that the CYP71A10 enzyme functions primarily as an N-demethylase with respect to fluometuron, linuron and diuron, with some N-demethoxylase activity also observed with linuron. Using chlortoluron as a substrate, the enzyme apparently functions primarily as a methyl-ring hydroxylase and to a lesser extent as an N-demethylase.

## EXAMPLE 6

### Herbicide Metabolism in Transgenic Tobacco

To determine whether overexpression of the soybean CYP71A10 cDNA

in a higher plant system enhances metabolism of phenylurea herbicides, the GUS gene in the binary vector pBI121 was excised and replaced with the CYP71A10 reading frame. This construct placed the CYP71A10 cDNA under the transcriptional control of the constitutive 35S promoter of Cauliflower Mosaic Virus; kanamycin selection was facilitated via the nptII selectable marker. Agrobacterium-mediated transformation of *Nicotiana tabacum* cv SR1 leaf discs resulted in the recovery of several dozen independent kanamycin-resistant transformants. The plants were subsequently grown to maturity in a greenhouse and allowed to set seed.

For the herbicide metabolism assays, seeds from one randomly selected transgenic line, designated 25/2, were germinated on kanamycin-containing media to eliminate potential nontransgenic segregants. Of 17 germinated seedlings grown, only one individual was inhibited by kanamycin (data not shown). This result suggests that line 25/2 possesses more than one independently segregating transgene. Individual leaves from the 25/2 progeny were excised and incubated with radiolabeled phenylurea herbicides. As shown in Table 5, leaves of the kanamycin-resistant individuals of line 25/2 metabolized all of the four herbicides tested to a much greater extent than the pBI121-transformed control plants.

The relative migrations of the metabolic products revealed by TLC suggest that the products observed in the *in vivo* excised leaf assay are primarily the same as were generated from the *in vitro* assays using yeast microsomes for fluometuron, linuron and diuron (data not shown). For chlortoluron, additional metabolites were also observed. These likely represent combinations of ring-methyl hydroxylated and mono- and di-demethylated species as had been observed by Shiota et al. *Pestic. Biochem. Physiol.* 54:190-198 (1996), in their analysis of chlortoluron-resistant transgenic tobacco that overexpressed the rat CYP1A1 gene. Differences in the ratios of the observed chlortoluron metabolites were also observed between the CYP71A10-transformed and the control plants.

Sixty three percent of the metabolites produced in the control leaves was N-

demethyl chlortoluron; in contrast, ring-methyl hydroxy chlortoluron was the most abundant metabolite generated in the CYP71A10-transformed leaves (47%) and only 8% of the metabolites represented N-demethyl chlortoluron.

5

Table 5

**Phenylurea Metabolism after 14 Hours by Excised Leaves of Transgenic Tobacco Plant 25/2 Progeny**

Herbicide <sup>a</sup>	CYP71A10-transformed	Control <sup>b</sup>
	% of herbicide metabolized	
Fluometuron	91 (4.5) <sup>c</sup>	15 (0.6)
Linuron	87 (2.0)	12 (2.6)
Chlortoluron	85 (8.1) <sup>d</sup>	39 (7.5) <sup>d</sup>
Diuron	49 (7.0)	20 (2.0)

- (a) Equal amounts of herbicide (1.2 nmol) were added for each experiment.
- (b) Plants transformed with the pBI121 construct were used as controls.
- (c) Values in parentheses represent standard error. A single leaf was assayed from four independent 25/2 plants and three independent control plants.
- (d) The major chlortoluron metabolite in the control plants represented N-demethyl chlortoluron (63%). The metabolites recovered from the CYP71A10-transformed leaves were ring-methyl hydroxy chlortoluron (47%), N-demethyl chlortoluron (8%) and other derivatives (45%).

25

**EXAMPLE 7****Herbicide Tolerance**

To establish whether enhanced herbicide metabolism leads to an increase in tolerance at the whole plant level, seeds from transgenic plant 25/2 were germinated on an agarose-base medium containing MS salts and varying

concentrations of linuron. Growth of wild-type SR1 plants and transgenic control plants expressing the GUS gene (from vector pBI121) was severely inhibited when exposed to 0.25  $\mu$ M linuron and completely arrested at concentrations of 0.5  $\mu$ M and higher (data not shown). As shown in Figure 5, progeny of plant 5 25/2 grown on media containing no herbicide (Figure 5A) appeared indistinguishable from the same seed grown in the presence of 0.5  $\mu$ M linuron (Figure 5C), where only one of 23 germinated seedlings appeared to be inhibited by the herbicide. This ratio appears to be consistent with that observed when seeds from the same parent were grown on selective media containing 10 kanamycin; only one of 17 seedlings failed to grow in the presence of kanamycin. Figure 5B shows control tobacco plants (transformed with vector pBI121), grown on media containing 0.5  $\mu$ M linuron. 25/2 plants tolerant to linuron levels as high as 2.5  $\mu$ M linuron were observed, although an increasing percentage of the plants showed growth inhibition as the herbicide concentration 15 was increased (Figure 5D). Segregation of the transgene(s) may be leading to variability in expression levels among the progeny of 25/2.

To examine whether the acquisition of herbicide tolerance is unique to line 25/2, seeds from 20 other independent CYP71A10-expressing transgenic plants were similarly germinated and grown on media containing 0.5  $\mu$ M 20 linuron. Of these, 19 lines gave rise to progeny that were linuron tolerant. The percentage of tolerant individuals for each line varied from approximately 20% to 100% (data not shown). This variation likely represents differences in the copy number, expression levels and segregation of the transgene among the independent lines.

25 Chlortoluron-tolerance of line 25/2 was also evident. At 1.0  $\mu$ M herbicide concentration chlortoluron completely arrested the growth of the control plants (Figure 5E). Although growth of the 25/2 plants was modestly inhibited at this herbicide concentration, with the exception of two presumably nontransgenic segregants, the CYP71A10-transformed plants appeared healthy 30 (Figure 5F). In contrast to linuron and chlortoluron, little tolerance of line 25/2

-38-

to fluometuron or diuron was observed. Herbicide concentrations that were injurious to the control plants also inhibited the growth of line 25/2 individuals. Enhanced fluometuron or diuron tolerance was only observed at the very lowest herbicide concentrations necessary to impose growth inhibition in the control plants (data not shown).

5



-39-  
SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Siminszky, Balazs  
Dewey, Ralph E.  
Corbin, Frederick T.
- (ii) TITLE OF INVENTION: Novel Cytochrome P-450 Constructs and  
Methods of Producing Herbicide-Resistant Transgenic Plants
- (iii) NUMBER OF SEQUENCES: 23
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Virginia C. Bennett
  - (B) STREET: PO Box 37428
  - (C) CITY: Raleigh
  - (D) STATE: North Carolina
  - (E) COUNTRY: USA
  - (F) ZIP: 27627
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Bennett, Virginia C.
  - (B) REGISTRATION NUMBER: 37,092
  - (C) REFERENCE/DOCKET NUMBER: 5051-409
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 919-854-1400
  - (B) TELEFAX: 919-854-1401

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1838 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 4..1542

-40-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AAA ATG GCT CTA CTA TCA TCA GTC CTA AAG CAA TTG CCG CAT GAG CTA	48
Met Ala Leu Leu Ser Ser Val Leu Lys Gln Leu Pro His Glu Leu	
1 5 10 15	
AGT TCA ACC CAT TAC CTA ACA GTT TTC TTC TGC ATC TTC CTT ATA CTT	96
Ser Ser Thr His Tyr Leu Thr Val Phe Phe Cys Ile Phe Leu Ile Leu	
20 25 30	
CTT CAG CTA ATA AGA AGA AAC AAA TAC AAT CTG CCA CCA TCC CCA CCA	144
Leu Gln Leu Ile Arg Arg Asn Lys Tyr Asn Leu Pro Pro Ser Pro Pro	
35 40 45	
AAG ATA CCC ATA ATC GGC AAT CTT CAC CAG CTA GGC ACA CTG CCA CAC	192
Lys Ile Pro Ile Ile Gly Asn Leu His Gln Leu Gly Thr Leu Pro His	
50 55 60	
CGC TCC TTT CAT GCA CTC TCA CAC AAA TAT GGC CCT CTC ATG ATG TTG	240
Arg Ser Phe His Ala Leu Ser His Lys Tyr Gly Pro Leu Met Met Leu	
65 70 75	
CAA TTG GGT CAA ATT CCA ACC CTA GTG GTC TCA TCA GCT GAC GTG GCC	288
Gln Leu Gly Gln Ile Pro Thr Leu Val Val Ser Ser Ala Asp Val Ala	
80 85 90 95	
AGA GAA ATA ATC AAA ACG CAT GAT GTT GTT TTC TCC AAC CGC CGA CAA	336
Arg Glu Ile Ile Lys Thr His Asp Val Val Phe Ser Asn Arg Arg Gln	
100 105 110	
CCT ACA GCT GCT AAA ATC TTT GGT TAT GGA TGC AAA GAT GTG GCT TTC	384
Pro Thr Ala Ala Lys Ile Phe Gly Tyr Gly Cys Lys Asp Val Ala Phe	
115 120 125	
GTG TAC TAC CGC GAA GAG TGG AGA CAA AAG ATA AAG ACA TGT AAG GTT	432
Val Tyr Tyr Arg Glu Glu Trp Arg Gln Lys Ile Lys Thr Cys Lys Val	
130 135 140	
GAG CTT ATG AGT CTG AAG AAG GTG CGG TTG TTT CAT TCC ATT AGA CAA	480
Glu Leu Met Ser Leu Lys Lys Val Arg Leu Phe His Ser Ile Arg Gln	
145 150 155	
GAA GTT GTT ACA GAG TTG GTT GAA GCT ATA GGT GAA GCG TGT GGT AGT	528
Glu Val Val Thr Glu Leu Val Glu Ala Ile Gly Glu Ala Cys Gly Ser	
160 165 170 175	
GAA AGA CCA TGT GTG AAT CTG ACT GAG ATG CTG ATG GCA GCA TCG AAC	576
Glu Arg Pro Cys Val Asn Leu Thr Glu Met Leu Met Ala Ala Ser Asn	
180 185 190	
GAC ATT GTG TCT AGA TGT GTT CTT GGA CGG AAG TGT GAT GAT GCA TGT	624
Asp Ile Val Ser Arg Cys Val Leu Gly Arg Lys Cys Asp Asp Ala Cys	
195 200 205	
GGT GGT AGT GGC AGT AGC AGC TTT GCA GCG TTG GGA AGA AAG ATT ATG	672
Gly Gly Ser Gly Ser Ser Ser Phe Ala Ala Leu Gly Arg Lys Ile Met	
210 215 220	
AGA CTA TTA TCG GCT TTC AGC GTG GGT GAT TTC TTC CCT TCG TTG GGT	720

-41-

Arg	Leu	Leu	Ser	Ala	Phe	Ser	Val	Gly	Asp	Phe	Phe	Pro	Ser	Leu	Gly	
225						230					235					
TGG	GTT	GAC	TAT	CTG	ACT	GGC	TTA	ATT	CCA	GAG	ATG	AAA	ACC	ACG	TTT	768
Trp	Val	Asp	Tyr	Leu	Thr	Gly	Leu	Ile	Pro	Glu	Met	Lys	Thr	Thr	Phe	
240					245					250					255	
CTC	GCA	GTA	GAT	GCT	TTC	CTT	GAT	GAG	GTA	ATT	GCA	GAA	CAC	GAG	AGC	816
Leu	Ala	Val	Asp	Ala	Phe	Leu	Asp	Glu	Val	Ile	Ala	Glu	His	Glu	Ser	
				260					265					270		
AGT	AAC	AAG	AAG	AAT	GAT	GAC	TTC	TTG	GGG	ATA	CTT	CTT	CAA	CTT	CAA	864
Ser	Asn	Lys	Lys	Asn	Asp	Asp	Phe	Leu	Gly	Ile	Leu	Leu	Gln	Leu	Gln	
			275					280					285			
GAA	TGT	GGG	AGG	CTT	GAC	TTT	CAG	CTC	GAC	CGA	GAT	AAC	CTC	AAA	GCA	912
Glu	Cys	Gly	Arg	Leu	Asp	Phe	Gln	Leu	Asp	Arg	Asp	Asn	Leu	Lys	Ala	
		290					295					300				
ATC	CTA	GTG	GAC	ATG	ATA	ATA	GGT	GGG	AGT	GAC	ACT	ACT	TCA	ACA	ACT	960
Ile	Leu	Val	Asp	Met	Ile	Ile	Gly	Gly	Ser	Asp	Thr	Thr	Ser	Thr	Thr	
	305					310					315					
CTA	GAA	TGG	ACT	TTT	GCG	GAG	TTC	CTT	AGA	AAT	CCA	AAT	ACC	ATG	AAG	1008
Leu	Glu	Trp	Thr	Phe	Ala	Glu	Phe	Leu	Arg	Asn	Pro	Asn	Thr	Met	Lys	
320					325					330					335	
AAA	GCT	CAA	GAA	GAG	GTA	AGA	AGA	GTG	GTG	GGA	ATC	AAT	TCC	AAA	GCA	1056
Lys	Ala	Gln	Glu	Glu	Val	Arg	Arg	Val	Val	Gly	Ile	Asn	Ser	Lys	Ala	
				340					345					350		
GTA	CTG	GAT	GAA	AAT	TGT	GTG	AAT	CAA	ATG	AAC	TAC	TTG	AAA	TGT	GTA	1104
Val	Leu	Asp	Glu	Asn	Cys	Val	Asn	Gln	Met	Asn	Tyr	Leu	Lys	Cys	Val	
			355					360					365			
GTC	AAA	GAA	ACT	TTG	AGA	TTA	CAT	CCA	CCC	CTT	CCT	CTT	TTG	ATT	GCT	1152
Val	Lys	Glu	Thr	Leu	Arg	Leu	His	Pro	Pro	Leu	Pro	Leu	Leu	Ile	Ala	
	370						375					380				
CGA	GAG	ACA	TCA	TCA	AGT	GTA	AAA	CTA	AGA	GGG	TAC	GAT	ATT	CCC	GCA	1200
Arg	Glu	Thr	Ser	Ser	Ser	Val	Lys	Leu	Arg	Gly	Tyr	Asp	Ile	Pro	Ala	
	385					390					395					
AAA	ACA	ATG	GTA	TTT	ATC	AAT	GCA	TGG	GCG	ATC	CAG	AGG	GAT	CCT	GAA	1248
Lys	Thr	Met	Val	Phe	Ile	Asn	Ala	Trp	Ala	Ile	Gln	Arg	Asp	Pro	Glu	
400					405					410					415	
TTA	TGG	GAT	GAT	CCT	GAA	GAA	TTT	ATT	CCC	GAA	AGA	TTT	GAA	ACT	AGC	1296
Leu	Trp	Asp	Asp	Pro	Glu	Glu	Phe	Ile	Pro	Glu	Arg	Phe	Glu	Thr	Ser	
				420					425					430		
CAA	GTT	GAT	CTT	AAT	GGA	CAA	GAT	TTT	CAA	TTA	ATT	CCG	TTC	GGT	ATT	1344
Gln	Val	Asp	Leu	Asn	Gly	Gln	Asp	Phe	Gln	Leu	Ile	Pro	Phe	Gly	Ile	
			435					440					445			
GGG	AGA	AGG	GGA	TGC	CCT	GCA	ATG	TCA	TTT	GGA	CTT	GCT	TCA	ACT	GAG	1392
Gly	Arg	Arg	Gly	Cys	Pro	Ala	Met	Ser	Phe	Gly	Leu	Ala	Ser	Thr	Glu	
	450						455					460				

-42-

TAT GTT CTT GCT AAT CTT TTG TAT TGG TTC AAT TGG AAT ATG TCC GAG 1440  
 Tyr Val Leu Ala Asn Leu Leu Tyr Trp Phe Asn Trp Asn Met Ser Glu  
 465 470 475

TCT GGA CGT ATA TTG ATG CAC AAC ATT GAC ATG AGT GAG ACA AAT GGA 1488  
 Ser Gly Arg Ile Leu Met His Asn Ile Asp Met Ser Glu Thr Asn Gly  
 480 485 490 495

CTC ACT GTC AGT AAG AAA GTA CCA CTT CAT CTT GAA CCA GAA CCA TAT 1536  
 Leu Thr Val Ser Lys Lys Val Pro Leu His Leu Glu Pro Glu Pro Tyr  
 500 505 510

AAA ACA TGATCATTTT ACATTATGCA TGTTTGGCAA CACCTATAAA GAGTATAGAT 1592  
 Lys Thr

CTGGAAGTAC TTCAATTTAG TAATGGATGT AAAAGCTATA CAATAAGAAG TGCTAACAAG 1652

CTAGGATATG AGCATTATG GAGTAACGAG TGAGGTTCCA AAGAGTCTAA TTACTCGTCT 1712

CTTGAACATT GTTATATTTG TTTTCTTGCA GTTTGTTAAT CTTTGAATA GTTGTTCAC 1772

ATTTATTTTT GTATGGTTTG TTGGTATGTT GTGGAAGGCG TTAGTAAAAA TTTGTGGTGT 1832

GTTCTT 1838

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 513 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Leu Leu Ser Ser Val Leu Lys Gln Leu Pro His Glu Leu Ser  
 1 5 10 15

Ser Thr His Tyr Leu Thr Val Phe Phe Cys Ile Phe Leu Ile Leu Leu  
 20 25 30

Gln Leu Ile Arg Arg Asn Lys Tyr Asn Leu Pro Pro Ser Pro Pro Lys  
 35 40 45

Ile Pro Ile Ile Gly Asn Leu His Gln Leu Gly Thr Leu Pro His Arg  
 50 55 60

Ser Phe His Ala Leu Ser His Lys Tyr Gly Pro Leu Met Met Leu Gln  
 65 70 75 80

Leu Gly Gln Ile Pro Thr Leu Val Val Ser Ser Ala Asp Val Ala Arg  
 85 90 95

Glu Ile Ile Lys Thr His Asp Val Val Phe Ser Asn Arg Arg Gln Pro  
 100 105 110

Thr Ala Ala Lys Ile Phe Gly Tyr Gly Cys Lys Asp Val Ala Phe Val

-43-

115		120		125
Tyr Tyr Arg Glu Glu Trp Arg Gln Lys Ile Lys Thr Cys Lys Val Glu				
130		135		140
Leu Met Ser Leu Lys Lys Val Arg Leu Phe His Ser Ile Arg Gln Glu				
145		150		155
Val Val Thr Glu Leu Val Glu Ala Ile Gly Glu Ala Cys Gly Ser Glu				
		165		170
Arg Pro Cys Val Asn Leu Thr Glu Met Leu Met Ala Ala Ser Asn Asp				
		180		185
Ile Val Ser Arg Cys Val Leu Gly Arg Lys Cys Asp Asp Ala Cys Gly				
		195		200
Gly Ser Gly Ser Ser Ser Phe Ala Ala Leu Gly Arg Lys Ile Met Arg				
		210		215
Leu Leu Ser Ala Phe Ser Val Gly Asp Phe Phe Pro Ser Leu Gly Trp				
		225		230
Val Asp Tyr Leu Thr Gly Leu Ile Pro Glu Met Lys Thr Thr Phe Leu				
		245		250
Ala Val Asp Ala Phe Leu Asp Glu Val Ile Ala Glu His Glu Ser Ser				
		260		265
Asn Lys Lys Asn Asp Asp Phe Leu Gly Ile Leu Leu Gln Leu Gln Glu				
		275		280
Cys Gly Arg Leu Asp Phe Gln Leu Asp Arg Asp Asn Leu Lys Ala Ile				
		290		295
Leu Val Asp Met Ile Ile Gly Gly Ser Asp Thr Thr Ser Thr Thr Leu				
		305		310
Glu Trp Thr Phe Ala Glu Phe Leu Arg Asn Pro Asn Thr Met Lys Lys				
		325		330
Ala Gln Glu Glu Val Arg Arg Val Val Gly Ile Asn Ser Lys Ala Val				
		340		345
Leu Asp Glu Asn Cys Val Asn Gln Met Asn Tyr Leu Lys Cys Val Val				
		355		360
Lys Glu Thr Leu Arg Leu His Pro Pro Leu Pro Leu Leu Ile Ala Arg				
		370		375
Glu Thr Ser Ser Ser Val Lys Leu Arg Gly Tyr Asp Ile Pro Ala Lys				
		385		390
Thr Met Val Phe Ile Asn Ala Trp Ala Ile Gln Arg Asp Pro Glu Leu				
		405		410
Trp Asp Asp Pro Glu Glu Phe Ile Pro Glu Arg Phe Glu Thr Ser Gln				
		420		425
Val Asp Leu Asn Gly Gln Asp Phe Gln Leu Ile Pro Phe Gly Ile Gly				

Thr

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1691 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: CDS  
(B) LOCATION: 16..1545

CCTAGATCTA	TCATC	ATG	GTC	ATG	GAG	CTT	CAC	AAC	CAC	ACC	CCT	TTC	TCT			51
		Met	Val	Met	Glu	Leu	His	Asn	His	Thr	Pro	Phe	Ser			
		1				5					10					
ATT	TAC	TTC	ATT	ACC	TCC	ATT	CTC	TTT	ATT	TTC	TTC	GTG	TTC	TTC	AAA	99
Ile	Tyr	Phe	Ile	Thr	Ser	Ile	Leu	Phe	Ile	Phe	Phe	Val	Phe	Phe	Lys	
		15				20						25				
TTA	GTT	CAA	AGA	TCG	GAT	TCC	AAA	ACC	TCC	TCT	ACC	TGC	AAA	TTG	CCC	147
Leu	Val	Gln	Arg	Ser	Asp	Ser	Lys	Thr	Ser	Ser	Thr	Cys	Lys	Leu	Pro	
		30				35					40					
CCA	GGA	CCA	AGG	ACA	CTA	CCT	CTC	ATA	GGG	AAC	ATA	CAC	CAG	ATT	GTT	195
Pro	Gly	Pro	Arg	Thr	Leu	Pro	Leu	Ile	Gly	Asn	Ile	His	Gln	Ile	Val	
		45			50					55					60	
GGC	TCA	CTG	CCG	GTT	CAT	TAC	TAC	TTA	AAA	AAT	TTG	GCA	GAT	AAG	TAT	243
Gly	Ser	Leu	Pro	Val	His	Tyr	Tyr	Leu	Lys	Asn	Leu	Ala	Asp	Lys	Tyr	
				65				70						75		
GGT	CCA	TTA	ATG	CAT	CTA	AAA	CTA	GGA	GAG	GTG	TCC	AAC	ATC	ATA	GTC	291
Gly	Pro	Leu	Met	His	Leu	Lys	Leu	Gly	Glu	Val	Ser	Asn	Ile	Ile	Val	
			80				85						90			
ACT	TCC	CCA	GAA	ATG	GCC	CAA	GAG	ATT	ATG	AAG	ACA	CAT	GAT	CTC	AAC	339

-45-

Thr Ser Pro Glu Met Ala Gln Glu Ile Met Lys Thr His Asp Leu Asn	
95 100 105	
TTC TCT GAT AGG CCA GAC TTT GTA TTG TCT AGA ATA GTT TCT TAC AAC	387
Phe Ser Asp Arg Pro Asp Phe Val Leu Ser Arg Ile Val Ser Tyr Asn	
110 115 120	
GGT TCT GGC ATT GTC TTC AGT CAA CAT GGA GAC TAT TGG AGG CAA CTA	435
Gly Ser Gly Ile Val Phe Ser Gln His Gly Asp Tyr Trp Arg Gln Leu	
125 130 135 140	
AGA AAG ATA TGC ACA GTA GAG TTA CTA ACA GCA AAG CGC GTG CAG TCT	483
Arg Lys Ile Cys Thr Val Glu Leu Leu Thr Ala Lys Arg Val Gln Ser	
145 150 155	
TTT CGG TCC ATA AGA GAA GAG GAG GTG GCA GAA CTA GTT AAA AAA ATA	531
Phe Arg Ser Ile Arg Glu Glu Glu Val Ala Glu Leu Val Lys Lys Ile	
160 165 170	
GCT GCA ACT GCA AGT GAA GAA GGG GGG TCC ATT TTT AAT CTC ACC CAG	579
Ala Ala Thr Ala Ser Glu Glu Gly Gly Ser Ile Phe Asn Leu Thr Gln	
175 180 185	
AGC ATT TAC TCA ATG ACT TTT GGG ATA GCG GCA CGA GCG GCT TTT GGT	627
Ser Ile Tyr Ser Met Thr Phe Gly Ile Ala Ala Arg Ala Ala Phe Gly	
190 195 200	
AAA AAG AGC AGA TAC CAA CAA GTG TTC ATA TCA AAC ATG CAT AAA CAA	675
Lys Lys Ser Arg Tyr Gln Gln Val Phe Ile Ser Asn Met His Lys Gln	
205 210 215 220	
TTG ATG CTT CTG GGA GGG TTT TCT GTT GCT GAT CTC TAT CCT TCT AGT	723
Leu Met Leu Leu Gly Gly Phe Ser Val Ala Asp Leu Tyr Pro Ser Ser	
225 230 235	
AGA GTG TTT CAA ATG ATG GGG GCG ACG GGG AAA CTT GAA AAA GTG CAT	771
Arg Val Phe Gln Met Met Gly Ala Thr Gly Lys Leu Glu Lys Val His	
240 245 250	
AGA GTG ACA GAT AGG GTG TTG CAA GAC ATC ATC GAC GAG CAC AAA AAT	819
Arg Val Thr Asp Arg Val Leu Gln Asp Ile Ile Asp Glu His Lys Asn	
255 260 265	
AGA AAC AGA AGC AGC GAG GAG CGT GAA GCA GTG GAA GAT CTA GTT GAT	867
Arg Asn Arg Ser Ser Glu Glu Arg Glu Ala Val Glu Asp Leu Val Asp	
270 275 280	
GTT CTT CTC AAG TTT CAA AAG GAA TCG GAA TTT CGC TTG ACT GAT GAC	915
Val Leu Leu Lys Phe Gln Lys Glu Ser Glu Phe Arg Leu Thr Asp Asp	
285 290 295 300	
AAC ATT AAA GCC GTC ATC CAG GAC ATA TTC ATT GGT GGA GGC GAA ACA	963
Asn Ile Lys Ala Val Ile Gln Asp Ile Phe Ile Gly Gly Gly Glu Thr	
305 310 315	
TCA TCT TCT GTT GTG GAA TGG GGG ATG TCA GAA TTG ATA AGA AAC CCG	1011
Ser Ser Ser Val Val Glu Trp Gly Met Ser Glu Leu Ile Arg Asn Pro	
320 325 330	

-46-

AGG GTG ATG GAA GAA GCA CAA GCA GAG GTG AGA AGA GTG TAT GAT AGC	1059
Arg Val Met Glu Glu Ala Gln Ala Glu Val Arg Arg Val Tyr Asp Ser	
335 340 345	
AAG GGA TAT GTG GAT GAG ACA GAA TTG CAC CAA TTG ATA TAC TTA AAG	1107
Lys Gly Tyr Val Asp Glu Thr Glu Leu His Gln Leu Ile Tyr Leu Lys	
350 355 360	
TCC ATC ATC AAA GAA ACC ATG AGG TTA CAT CCA CCT GTG CCA TTG TTA	1155
Ser Ile Ile Lys Glu Thr Met Arg Leu His Pro Pro Val Pro Leu Leu	
365 370 375 380	
GTT CCT AGA GTA AGT AGA GAA AGG TGC CAA ATC AAT GGA TAT GAG ATA	1203
Val Pro Arg Val Ser Arg Glu Arg Cys Gln Ile Asn Gly Tyr Glu Ile	
385 390 395	
CCC TCT AAG ACT AGG ATC ATT ATC AAT GCT TGG GCA ATT GGA AGG AAT	1251
Pro Ser Lys Thr Arg Ile Ile Ile Asn Ala Trp Ala Ile Gly Arg Asn	
400 405 410	
CCT AAG TAT TGG GGT GAA ACT GAG AGT TTT AAA CCT GAG AGG TTT CTT	1299
Pro Lys Tyr Trp Gly Glu Thr Glu Ser Phe Lys Pro Glu Arg Phe Leu	
415 420 425	
AAT AGC TCC ATT GAT TTT AGG GGC ACA GAC TTT GAA TTT ATC CCA TTT	1347
Asn Ser Ser Ile Asp Phe Arg Gly Thr Asp Phe Glu Phe Ile Pro Phe	
430 435 440	
GGT GCT GGA AGG AGG ATC TGC CCC GGC ATT ACA TTT GCC ATA CCC AAC	1395
Gly Ala Gly Arg Arg Ile Cys Pro Gly Ile Thr Phe Ala Ile Pro Asn	
445 450 455 460	
ATT GAG TTG CCA CTT GCT CAG TTA CTT TAC CAC TTT GAT TGG AAG CTT	1443
Ile Glu Leu Pro Leu Ala Gln Leu Leu Tyr His Phe Asp Trp Lys Leu	
465 470 475	
CCC AAT AAA ATG AAG AAT GAA GAA CTT GAC ATG ACG GAG TCA AAT GGA	1491
Pro Asn Lys Met Lys Asn Glu Glu Leu Asp Met Thr Glu Ser Asn Gly	
480 485 490	
ATT ACT TTA CGA AGA CAA AAT GAC CTC TGC TTG ATT CCC ATT ACT CGT	1539
Ile Thr Leu Arg Arg Gln Asn Asp Leu Cys Leu Ile Pro Ile Thr Arg	
495 500 505	
CTA CCT TAAAAATGTAT GAACAATTAA TGTCATAAAC TATTTAAGTT TTATCTTTTA	1595
Leu Pro	
510	
CTACTTCCAG CATTTTCGTAA TTGGACAATG ACTATGATTA ACTTAAGTTA CTTCCATTATG	1655
ATTAACCTTGA CATATGAATG AACATTTCTA AGATAA	1691

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 510 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear



-47-

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Val Met Glu Leu His Asn His Thr Pro Phe Ser Ile Tyr Phe Ile
 1             5             10             15

Thr Ser Ile Leu Phe Ile Phe Phe Val Phe Phe Lys Leu Val Gln Arg
 20             25             30

Ser Asp Ser Lys Thr Ser Ser Thr Cys Lys Leu Pro Pro Gly Pro Arg
 35             40             45

Thr Leu Pro Leu Ile Gly Asn Ile His Gln Ile Val Gly Ser Leu Pro
 50             55             60

Val His Tyr Tyr Leu Lys Asn Leu Ala Asp Lys Tyr Gly Pro Leu Met
 65             70             75             80

His Leu Lys Leu Gly Glu Val Ser Asn Ile Ile Val Thr Ser Pro Glu
 85             90             95

Met Ala Gln Glu Ile Met Lys Thr His Asp Leu Asn Phe Ser Asp Arg
100             105             110

Pro Asp Phe Val Leu Ser Arg Ile Val Ser Tyr Asn Gly Ser Gly Ile
115             120             125

Val Phe Ser Gln His Gly Asp Tyr Trp Arg Gln Leu Arg Lys Ile Cys
130             135             140

Thr Val Glu Leu Leu Thr Ala Lys Arg Val Gln Ser Phe Arg Ser Ile
145             150             155             160

Arg Glu Glu Glu Val Ala Glu Leu Val Lys Lys Ile Ala Ala Thr Ala
165             170             175

Ser Glu Glu Gly Gly Ser Ile Phe Asn Leu Thr Gln Ser Ile Tyr Ser
180             185             190

Met Thr Phe Gly Ile Ala Ala Arg Ala Ala Phe Gly Lys Lys Ser Arg
195             200             205

Tyr Gln Gln Val Phe Ile Ser Asn Met His Lys Gln Leu Met Leu Leu
210             215             220

Gly Gly Phe Ser Val Ala Asp Leu Tyr Pro Ser Ser Arg Val Phe Gln
225             230             235             240

Met Met Gly Ala Thr Gly Lys Leu Glu Lys Val His Arg Val Thr Asp
245             250             255

Arg Val Leu Gln Asp Ile Ile Asp Glu His Lys Asn Arg Asn Arg Ser
260             265             270

Ser Glu Glu Arg Glu Ala Val Glu Asp Leu Val Asp Val Leu Leu Lys
275             280             285

Phe Gln Lys Glu Ser Glu Phe Arg Leu Thr Asp Asp Asn Ile Lys Ala

```

-48-

```

      290              295              300
Val Ile Gln Asp Ile Phe Ile Gly Gly Gly Glu Thr Ser Ser Ser Val
305              310              315              320

Val Glu Trp Gly Met Ser Glu Leu Ile Arg Asn Pro Arg Val Met Glu
              325              330              335

Glu Ala Gln Ala Glu Val Arg Arg Val Tyr Asp Ser Lys Gly Tyr Val
              340              345              350

Asp Glu Thr Glu Leu His Gln Leu Ile Tyr Leu Lys Ser Ile Ile Lys
              355              360              365

Glu Thr Met Arg Leu His Pro Pro Val Pro Leu Leu Val Pro Arg Val
              370              375              380

Ser Arg Glu Arg Cys Gln Ile Asn Gly Tyr Glu Ile Pro Ser Lys Thr
385              390              395              400

Arg Ile Ile Ile Asn Ala Trp Ala Ile Gly Arg Asn Pro Lys Tyr Trp
              405              410              415

Gly Glu Thr Glu Ser Phe Lys Pro Glu Arg Phe Leu Asn Ser Ser Ile
              420              425              430

Asp Phe Arg Gly Thr Asp Phe Glu Phe Ile Pro Phe Gly Ala Gly Arg
              435              440              445

Arg Ile Cys Pro Gly Ile Thr Phe Ala Ile Pro Asn Ile Glu Leu Pro
              450              455              460

Leu Ala Gln Leu Leu Tyr His Phe Asp Trp Lys Leu Pro Asn Lys Met
465              470              475              480

Lys Asn Glu Glu Leu Asp Met Thr Glu Ser Asn Gly Ile Thr Leu Arg
              485              490              495

Arg Gln Asn Asp Leu Cys Leu Ile Pro Ile Thr Arg Leu Pro
              500              505              510

```

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1644 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 4..1542

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AAA ATG GCC ACT CTT TCC TCC TAC GAC CAC TTC ATC TTC ACT GCC TTA

48

-49-

Met	Ala	Thr	Leu	Ser	Ser	Tyr	Asp	His	Phe	Ile	Phe	Thr	Ala	Leu		
1				5					10					15		
GCT	TTC	TTC	ATA	TCT	GGC	CTA	ATT	TTC	TTC	CTC	AAA	CAG	AAA	TCC	AAA	96
Ala	Phe	Phe	Ile	Ser	Gly	Leu	Ile	Phe	Phe	Leu	Lys	Gln	Lys	Ser	Lys	
			20					25					30			
TCC	AAA	AAG	TTC	AAC	CTC	CCT	CCA	GGA	CCC	CCC	GGG	TGG	CCT	ATT	GTT	144
Ser	Lys	Lys	Phe	Asn	Leu	Pro	Pro	Gly	Pro	Pro	Gly	Trp	Pro	Ile	Val	
			35					40					45			
GGG	AAC	CTC	TTC	CAA	GTT	GCT	CGT	TCT	GGG	AAA	CCT	TTC	TTT	GAG	TAT	192
Gly	Asn	Leu	Phe	Gln	Val	Ala	Arg	Ser	Gly	Lys	Pro	Phe	Phe	Glu	Tyr	
	50					55					60					
GTG	AAC	GAT	GTG	AGA	CTC	AAA	TAT	GGC	TCA	ATC	TTC	ACC	CTC	AAG	ATG	240
Val	Asn	Asp	Val	Arg	Leu	Lys	Tyr	Gly	Ser	Ile	Phe	Thr	Leu	Lys	Met	
	65					70					75					
GGA	ACA	AGG	ACC	ATG	ATC	ATC	CTC	ACC	GAC	GCA	AAA	CTG	GTC	CAC	GAG	288
Gly	Thr	Arg	Thr	Met	Ile	Ile	Leu	Thr	Asp	Ala	Lys	Leu	Val	His	Glu	
	80				85					90					95	
GCC	ATG	ATC	CAA	AAG	GGT	GCA	ACC	TAC	GCC	ACC	AGG	CCC	CCC	GAG	AAC	336
Ala	Met	Ile	Gln	Lys	Gly	Ala	Thr	Tyr	Ala	Thr	Arg	Pro	Pro	Glu	Asn	
			100					105						110		
CCC	ACC	AGA	ACC	ATC	TTC	AGT	GAA	AAC	AAG	TTC	ACC	GTG	AAT	GCA	GCG	384
Pro	Thr	Arg	Thr	Ile	Phe	Ser	Glu	Asn	Lys	Phe	Thr	Val	Asn	Ala	Ala	
			115					120					125			
ACC	TAT	GGC	CCC	GTG	TGG	AAG	TCG	CTG	AGG	AGG	AAC	ATG	GTG	CAG	AAC	432
Thr	Tyr	Gly	Pro	Val	Trp	Lys	Ser	Leu	Arg	Arg	Asn	Met	Val	Gln	Asn	
	130					135						140				
ATG	CTC	AGC	TCA	ACA	AGA	CTT	AAG	GAG	TTT	CGC	AGT	GTT	CGG	GAC	AAT	480
Met	Leu	Ser	Ser	Thr	Arg	Leu	Lys	Glu	Phe	Arg	Ser	Val	Arg	Asp	Asn	
	145					150					155					
GCG	ATG	GAC	AAG	CTC	ATC	AAC	AGA	CTC	AAG	GAC	GAG	GCC	GAG	AAG	AAT	528
Ala	Met	Asp	Lys	Leu	Ile	Asn	Arg	Leu	Lys	Asp	Glu	Ala	Glu	Lys	Asn	
	160				165					170				175		
AAC	GGC	GTG	GTT	TGG	GTG	CTC	AAG	GAT	GCC	AGG	TTT	GCT	GTT	TTT	TGC	576
Asn	Gly	Val	Val	Trp	Val	Leu	Lys	Asp	Ala	Arg	Phe	Ala	Val	Phe	Cys	
			180					185					190			
ATA	CTT	GTG	GCT	ATG	TGT	TTT	GGT	CTT	GAG	ATG	GAT	GAG	GAG	ACA	GTG	624
Ile	Leu	Val	Ala	Met	Cys	Phe	Gly	Leu	Glu	Met	Asp	Glu	Glu	Thr	Val	
			195					200					205			
GAG	AGA	ATA	GAT	CAG	GTT	ATG	AAG	AGT	GTT	CTC	ATC	ACT	TTG	GAC	CCG	672
Glu	Arg	Ile	Asp	Gln	Val	Met	Lys	Ser	Val	Leu	Ile	Thr	Leu	Asp	Pro	
	210						215					220				
AGA	ATT	GAT	GAC	TAT	CTT	CCA	ATT	CTA	AGC	CCC	TTT	TTC	TCA	AAG	CAA	720
Arg	Ile	Asp	Asp	Tyr	Leu	Pro	Ile	Leu	Ser	Pro	Phe	Phe	Ser	Lys	Gln	
	225					230					235					

-50-

AGA AAG AAA GCC TTG GAG GTT CGC AGA GAA CAG GTT GAG TTC TTA GTT	768
Arg Lys Lys Ala Leu Glu Val Arg Arg Glu Gln Val Glu Phe Leu Val	
240 245 250 255	
CCA ATT ATA GAA CAA AGA AGA AGA GCA ATT CAA AAC CCT GGG TCA GAT	816
Pro Ile Ile Glu Gln Arg Arg Arg Ala Ile Gln Asn Pro Gly Ser Asp	
260 265 270	
CAC ACC GCC ACA ACG TTT TCC TAC CTA GAC ACA CTT TTT GAC CTC AAA	864
His Thr Ala Thr Thr Phe Ser Tyr Leu Asp Thr Leu Phe Asp Leu Lys	
275 280 285	
GTT GAA GGG AAG AAA TCA GCA CCC TCT GAT GCA GAA TTG GTG TCT TTA	912
Val Glu Gly Lys Lys Ser Ala Pro Ser Asp Ala Glu Leu Val Ser Leu	
290 295 300	
TGC TCA GAG TTT CTT AAC GGT GGC ACA GAC ACA ACA GCA ACA GCG GTT	960
Cys Ser Glu Phe Leu Asn Gly Gly Thr Asp Thr Thr Ala Thr Ala Val	
305 310 315	
GAG TGG GGC ATA GCA CAG CTC ATA GCG AAC CCT AAC GTT CAG ACA AAG	1008
Glu Trp Gly Ile Ala Gln Leu Ile Ala Asn Pro Asn Val Gln Thr Lys	
320 325 330 335	
CTG TAC GAG GAA ATA AAG AGA ACG GTG GGA GAG AAG AAG GTG GAT GAA	1056
Leu Tyr Glu Glu Ile Lys Arg Thr Val Gly Glu Lys Lys Val Asp Glu	
340 345 350	
AAG GAC GTT GAG AAA ATG CCA TAC CTA CAC GCT GTG GTG AAG GAG CTT	1104
Lys Asp Val Glu Lys Met Pro Tyr Leu His Ala Val Val Lys Glu Leu	
355 360 365	
CTA AGA AAG CAC CCT CCA ACA CAC TTT GTG CTA ACA CAT GCT GTG ACT	1152
Leu Arg Lys His Pro Pro Thr His Phe Val Leu Thr His Ala Val Thr	
370 375 380	
GAG CCC ACC ACT TTG GGA GGG TAT GAC ATA CCA ATT GAT GCA AAT GTT	1200
Glu Pro Thr Thr Leu Gly Gly Tyr Asp Ile Pro Ile Asp Ala Asn Val	
385 390 395	
GAG GTG TAC ACA CCA GCC ATT GCT GAG GAC CCC AAA AAT TGG TTA AAC	1248
Glu Val Tyr Thr Pro Ala Ile Ala Glu Asp Pro Lys Asn Trp Leu Asn	
400 405 410 415	
CCT GAG AAG TTT GAC CCT GAG AGA TTC ATC TCT GGG GGT GAG GAA GCA	1296
Pro Glu Lys Phe Asp Pro Glu Arg Phe Ile Ser Gly Gly Glu Ala	
420 425 430	
GAC ATA ACT GGG GTC ACA GGG GTG AAG ATG ATG CCA TTT GGG GTT GGG	1344
Asp Ile Thr Gly Val Thr Gly Val Lys Met Met Pro Phe Gly Val Gly	
435 440 445	
AGA AGG ATT TGC CCT GGC TTG GCT ATG GCC ACA GTG CAT ATT CAC CTC	1392
Arg Arg Ile Cys Pro Gly Leu Ala Met Ala Thr Val His Ile His Leu	
450 455 460	
ATG ATG GCA AGG ATG GTG CAG GAG TTT GAG TGG GGT GCA TAC CCT CCA	1440
Met Met Ala Arg Met Val Gln Glu Phe Glu Trp Gly Ala Tyr Pro Pro	
465 470 475	

-51-

GAG AAG AAG ATG GAT TTC ACT GGC AAG TGG GAG TTC ACT GTG GTC ATG 1488  
 Glu Lys Lys Met Asp Phe Thr Gly Lys Trp Glu Phe Thr Val Val Met  
 480 485 490 495

AAG GAG TCT CTA AGA GCA ACC ATC AAA CCA AGA GGA GGA GAA AAA GTG 1536  
 Lys Glu Ser Leu Arg Ala Thr Ile Lys Pro Arg Gly Gly Glu Lys Val  
 500 505 510

AAG TTG TAAAATTTTC CTGCTTCTAT TCTTCTGGGT TTAAATTTTC ACAGACAACA 1592  
 Lys Leu

TAAATATTAT TGCTATTATC ATCATCATAT ATGTATACAT CATCATGGTT AC 1644

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 513 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Thr Leu Ser Ser Tyr Asp His Phe Ile Phe Thr Ala Leu Ala  
 1 5 10 15  
 Phe Phe Ile Ser Gly Leu Ile Phe Phe Leu Lys Gln Lys Ser Lys Ser  
 20 25 30  
 Lys Lys Phe Asn Leu Pro Pro Gly Pro Pro Gly Trp Pro Ile Val Gly  
 35 40 45  
 Asn Leu Phe Gln Val Ala Arg Ser Gly Lys Pro Phe Phe Glu Tyr Val  
 50 55 60  
 Asn Asp Val Arg Leu Lys Tyr Gly Ser Ile Phe Thr Leu Lys Met Gly  
 65 70 75 80  
 Thr Arg Thr Met Ile Ile Leu Thr Asp Ala Lys Leu Val His Glu Ala  
 85 90 95  
 Met Ile Gln Lys Gly Ala Thr Tyr Ala Thr Arg Pro Pro Glu Asn Pro  
 100 105 110  
 Thr Arg Thr Ile Phe Ser Glu Asn Lys Phe Thr Val Asn Ala Ala Thr  
 115 120 125  
 Tyr Gly Pro Val Trp Lys Ser Leu Arg Arg Asn Met Val Gln Asn Met  
 130 135 140  
 Leu Ser Ser Thr Arg Leu Lys Glu Phe Arg Ser Val Arg Asp Asn Ala  
 145 150 155 160  
 Met Asp Lys Leu Ile Asn Arg Leu Lys Asp Glu Ala Glu Lys Asn Asn  
 165 170 175

-52-

Gly Val Val Trp Val Leu Lys Asp Ala Arg Phe Ala Val Phe Cys Ile  
 180 185 190

Leu Val Ala Met Cys Phe Gly Leu Glu Met Asp Glu Glu Thr Val Glu  
 195 200 205

Arg Ile Asp Gln Val Met Lys Ser Val Leu Ile Thr Leu Asp Pro Arg  
 210 215 220

Ile Asp Asp Tyr Leu Pro Ile Leu Ser Pro Phe Phe Ser Lys Gln Arg  
 225 230 235 240

Lys Lys Ala Leu Glu Val Arg Arg Glu Gln Val Glu Phe Leu Val Pro  
 245 250 255

Ile Ile Glu Gln Arg Arg Arg Ala Ile Gln Asn Pro Gly Ser Asp His  
 260 265 270

Thr Ala Thr Thr Phe Ser Tyr Leu Asp Thr Leu Phe Asp Leu Lys Val  
 275 280 285

Glu Gly Lys Lys Ser Ala Pro Ser Asp Ala Glu Leu Val Ser Leu Cys  
 290 295 300

Ser Glu Phe Leu Asn Gly Gly Thr Asp Thr Thr Ala Thr Ala Val Glu  
 305 310 315 320

Trp Gly Ile Ala Gln Leu Ile Ala Asn Pro Asn Val Gln Thr Lys Leu  
 325 330 335

Tyr Glu Glu Ile Lys Arg Thr Val Gly Glu Lys Lys Val Asp Glu Lys  
 340 345 350

Asp Val Glu Lys Met Pro Tyr Leu His Ala Val Val Lys Glu Leu Leu  
 355 360 365

Arg Lys His Pro Pro Thr His Phe Val Leu Thr His Ala Val Thr Glu  
 370 375 380

Pro Thr Thr Leu Gly Gly Tyr Asp Ile Pro Ile Asp Ala Asn Val Glu  
 385 390 395 400

Val Tyr Thr Pro Ala Ile Ala Glu Asp Pro Lys Asn Trp Leu Asn Pro  
 405 410 415

Glu Lys Phe Asp Pro Glu Arg Phe Ile Ser Gly Gly Glu Glu Ala Asp  
 420 425 430

Ile Thr Gly Val Thr Gly Val Lys Met Met Pro Phe Gly Val Gly Arg  
 435 440 445

Arg Ile Cys Pro Gly Leu Ala Met Ala Thr Val His Ile His Leu Met  
 450 455 460

Met Ala Arg Met Val Gln Glu Phe Glu Trp Gly Ala Tyr Pro Pro Glu  
 465 470 475 480

Lys Lys Met Asp Phe Thr Gly Lys Trp Glu Phe Thr Val Val Met Lys  
 485 490 495

-53-

Glu Ser Leu Arg Ala Thr Ile Lys Pro Arg Gly Gly Glu Lys Val Lys  
                   500                  505                  510

Leu

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1611 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 20..1588

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AAGCACTATC CCTCCCACC ATG ACA AGC CAC ATT GAC GAC AAC CTC TGG ATA	52
Met Thr Ser His Ile Asp Asp Asn Leu Trp Ile	
1                                  5                                  10	
ATA GCC CTG ACC TCG AAA TGC ACC CAA GAA AAC CTT GCA TGG GTC CTT	100
Ile Ala Leu Thr Ser Lys Cys Thr Gln Glu Asn Leu Ala Trp Val Leu	
15                                  20                                  25	
TTG ATC ATG GGC TCA CTC TGG TTA ACC ATG ACT TTC TAT TAC TGG TCA	148
Leu Ile Met Gly Ser Leu Trp Leu Thr Met Thr Phe Tyr Tyr Trp Ser	
30                                  35                                  40	
CAC CCC GGT GGT CCT GCC TGG GGC AAG TAC TAC ACC TAC TCT CCC CCC	196
His Pro Gly Gly Pro Ala Trp Gly Lys Tyr Tyr Thr Tyr Ser Pro Pro	
45                                  50                                  55	
CTT TCA ATC ATT CCC GGT CCC AAA GGC TTC CCT CTT ATT GGA AGC ATG	244
Leu Ser Ile Ile Pro Gly Pro Lys Gly Phe Pro Leu Ile Gly Ser Met	
60                                  65                                  70                                  75	
GGC CTC ATG ACT TCC CTG GCC CAT CAC CGT ATC GCA GCC GCG GCC GCC	292
Gly Leu Met Thr Ser Leu Ala His His Arg Ile Ala Ala Ala Ala Ala	
80                                  85                                  90	
ACA TGC AGA GCC AAG CGC CTC ATG GCC TTT AGT CTC GGC GAC ACA CGT	340
Thr Cys Arg Ala Lys Arg Leu Met Ala Phe Ser Leu Gly Asp Thr Arg	
95                                  100                                  105	
GTC ATC GTC ACG TGC CAC CCC GAC GTG GCC AAG GAG ATT CTC AAC AGC	388
Val Ile Val Thr Cys His Pro Asp Val Ala Lys Glu Ile Leu Asn Ser	
110                                  115                                  120	
TCC GTC TTC GCC GAT CGT CCC GTC AAA GAA TCC GCA TAC AGC CTC ATG	436
Ser Val Phe Ala Asp Arg Pro Val Lys Glu Ser Ala Tyr Ser Leu Met	
125                                  130                                  135	

-54-

TTT AAC CGC GCC ATC GGC TTC GCC TCT TAC GGA GTT TAC TGG CGA AGC Phe Asn Arg Ala Ile Gly Phe Ala Ser Tyr Gly Val Tyr Trp Arg Ser 140 145 150 155	484
CTC AGG AGA ATC GCC TCT AAT CAC CTC TTC TGC CCC CGC CAG ATA AAA Leu Arg Arg Ile Ala Ser Asn His Leu Phe Cys Pro Arg Gln Ile Lys 160 165 170	532
GCC TCT GAG CTC CAA CGC TCT CAA ATC GCC GCC CAA ATG GTT CAC ATC Ala Ser Glu Leu Gln Arg Ser Gln Ile Ala Ala Gln Met Val His Ile 175 180 185	580
CTA AAT AAC AAG CGC CAC CGC AGC TTA CGT GTT CGC CAA GTG CTG AAA Leu Asn Asn Lys Arg His Arg Ser Leu Arg Val Arg Gln Val Leu Lys 190 195 200	628
AAG GCT TCG CTC AGT AAC ATG ATG TGC TCC GTG TTT GGA CAA GAG TAT Lys Ala Ser Leu Ser Asn Met Met Cys Ser Val Phe Gly Gln Glu Tyr 205 210 215	676
AAG CTG CAC GAC CCA AAC AGC GGA ATG GAA GAC CTT GGA ATA TTA GTG Lys Leu His Asp Pro Asn Ser Gly Met Glu Asp Leu Gly Ile Leu Val 220 225 230 235	724
GAC CAA GGT TAT GAC CTG TTG GGC CTG TTT AAT TGG GCC GAC CAC CTT Asp Gln Gly Tyr Asp Leu Leu Gly Leu Phe Asn Trp Ala Asp His Leu 240 245 250	772
CCT TTT CTT GCA CAT TTC GAC GCC CAA AAT ATC CGG TTC AGG TGC TCC Pro Phe Leu Ala His Phe Asp Ala Gln Asn Ile Arg Phe Arg Cys Ser 255 260 265	820
AAC CTC GTC CCC ATG GTG AAC CGT TTC GTC GGC ACA ATC ATC GCT GAA Asn Leu Val Pro Met Val Asn Arg Phe Val Gly Thr Ile Ile Ala Glu 270 275 280	868
CAC CGA GCT AGT AAA ACC GAA ACC AAT CGT GAT TTT GTT GAC GTC TTG His Arg Ala Ser Lys Thr Glu Thr Asn Arg Asp Phe Val Asp Val Leu 285 290 295	916
CTC TCT CTC CCG GAA CCT GAT CAA TTA TCA GAC TCC GAC ATG ATC GCT Leu Ser Leu Pro Glu Pro Asp Gln Leu Ser Asp Ser Asp Met Ile Ala 300 305 310 315	964
GTA CTT TGG GAA ATG ATA TTC AGA GGA ACG GAC ACG GTA GCG GTT TTG Val Leu Trp Glu Met Ile Phe Arg Gly Thr Asp Thr Val Ala Val Leu 320 325 330	1012
ATA GAG TGG ATA CTC GCG AGG ATG GCG CTT CAT CCT CAT GTG CAG TCC Ile Glu Trp Ile Leu Ala Arg Met Ala Leu His Pro His Val Gln Ser 335 340 345	1060
AAA GTT CAA GAG GAG CTA GAT GCA GTT GTC GGA AAA GCA CGC GCC GTC Lys Val Gln Glu Glu Leu Asp Ala Val Val Gly Lys Ala Arg Ala Val 350 355 360	1108
GCA GAG GAT GAC GTG GCA GTG ATG ACG TAC CTA CCA GCG GTG GTG AAG Ala Glu Asp Asp Val Ala Val Met Thr Tyr Leu Pro Ala Val Val Lys 365 370 375	1156



-55-

GAG GTG CTG CGG CTG CAC CCG CCG GGC CCA CTT CTA TCA TGG GCC CGC	1204
Glu Val Leu Arg Leu His Pro Pro Gly Pro Leu Leu Ser Trp Ala Arg	
380 385 390 395	
TTG TCC ATC AAT GAT ACG ACC ATT GAT GGG TAT CAC GTA CCT GCG GGG	1252
Leu Ser Ile Asn Asp Thr Thr Ile Asp Gly Tyr His Val Pro Ala Gly	
400 405 410	
ACC ACT GCT ATG GTC AAC ACG TGG GCT ATT TGC AGG GAC CCA CAC GTG	1300
Thr Thr Ala Met Val Asn Thr Trp Ala Ile Cys Arg Asp Pro His Val	
415 420 425	
TGG AAG GAC CCA CTC GAA TTT ATG CCC GAG AGG TTT GTC ACT GCG GGT	1348
Trp Lys Asp Pro Leu Glu Phe Met Pro Glu Arg Phe Val Thr Ala Gly	
430 435 440	
GGA GAT GCC GAA TTT TCG ATA CTC GGG TCG GAT CCA AGA CTT GCT CCA	1396
Gly Asp Ala Glu Phe Ser Ile Leu Gly Ser Asp Pro Arg Leu Ala Pro	
445 450 455	
TTT GGG TCG GGT AGG AGA GCG TGC CCA GGG AAG ACT CTT GGA TGG GCT	1444
Phe Gly Ser Gly Arg Arg Ala Cys Pro Gly Lys Thr Leu Gly Trp Ala	
460 465 470 475	
ACG GTG AAC TTT TGG GTG GCG TCG CTC TTG CAT GAG TTC GAA TGG GTA	1492
Thr Val Asn Phe Trp Val Ala Ser Leu Leu His Glu Phe Glu Trp Val	
480 485 490	
CCG TCT GAT GAG AAG GGT GTT GAT CTG ACG GAG GTG CTG AAG CTC TCT	1540
Pro Ser Asp Glu Lys Gly Val Asp Leu Thr Glu Val Leu Lys Leu Ser	
495 500 505	
AGT GAA ATG GCT AAC CCT CTC ACC GTC AAA GTG CGC CCC AGG CGT GGA	1588
Ser Glu Met Ala Asn Pro Leu Thr Val Lys Val Arg Pro Arg Arg Gly	
510 515 520	
TAAGAGAGAG TTGAAGCTTT TAT	1611

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 523 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Thr Ser His Ile Asp Asp Asn Leu Trp Ile Ile Ala Leu Thr Ser	
1 5 10 15	
Lys Cys Thr Gln Glu Asn Leu Ala Trp Val Leu Leu Ile Met Gly Ser	
20 25 30	
Leu Trp Leu Thr Met Thr Phe Tyr Tyr Trp Ser His Pro Gly Gly Pro	
35 40 45	

-56-

Ala Trp Gly Lys Tyr Tyr Thr Tyr Ser Pro Pro Leu Ser Ile Ile Pro  
 50 55 60

Gly Pro Lys Gly Phe Pro Leu Ile Gly Ser Met Gly Leu Met Thr Ser  
 65 70 75 80

Leu Ala His His Arg Ile Ala Ala Ala Ala Thr Cys Arg Ala Lys  
 85 90 95

Arg Leu Met Ala Phe Ser Leu Gly Asp Thr Arg Val Ile Val Thr Cys  
 100 105 110

His Pro Asp Val Ala Lys Glu Ile Leu Asn Ser Ser Val Phe Ala Asp  
 115 120 125

Arg Pro Val Lys Glu Ser Ala Tyr Ser Leu Met Phe Asn Arg Ala Ile  
 130 135 140

Gly Phe Ala Ser Tyr Gly Val Tyr Trp Arg Ser Leu Arg Arg Ile Ala  
 145 150 155 160

Ser Asn His Leu Phe Cys Pro Arg Gln Ile Lys Ala Ser Glu Leu Gln  
 165 170 175

Arg Ser Gln Ile Ala Ala Gln Met Val His Ile Leu Asn Asn Lys Arg  
 180 185 190

His Arg Ser Leu Arg Val Arg Gln Val Leu Lys Lys Ala Ser Leu Ser  
 195 200 205

Asn Met Met Cys Ser Val Phe Gly Gln Glu Tyr Lys Leu His Asp Pro  
 210 215 220

Asn Ser Gly Met Glu Asp Leu Gly Ile Leu Val Asp Gln Gly Tyr Asp  
 225 230 235 240

Leu Leu Gly Leu Phe Asn Trp Ala Asp His Leu Pro Phe Leu Ala His  
 245 250 255

Phe Asp Ala Gln Asn Ile Arg Phe Arg Cys Ser Asn Leu Val Pro Met  
 260 265 270

Val Asn Arg Phe Val Gly Thr Ile Ile Ala Glu His Arg Ala Ser Lys  
 275 280 285

Thr Glu Thr Asn Arg Asp Phe Val Asp Val Leu Leu Ser Leu Pro Glu  
 290 295 300

Pro Asp Gln Leu Ser Asp Ser Asp Met Ile Ala Val Leu Trp Glu Met  
 305 310 315 320

Ile Phe Arg Gly Thr Asp Thr Val Ala Val Leu Ile Glu Trp Ile Leu  
 325 330 335

Ala Arg Met Ala Leu His Pro His Val Gln Ser Lys Val Gln Glu Glu  
 340 345 350

Leu Asp Ala Val Val Gly Lys Ala Arg Ala Val Ala Glu Asp Asp Val  
 355 360 365

-57-

Ala Val Met Thr Tyr Leu Pro Ala Val Val Lys Glu Val Leu Arg Leu  
 370 375 380

His Pro Pro Gly Pro Leu Leu Ser Trp Ala Arg Leu Ser Ile Asn Asp  
 385 390 395 400

Thr Thr Ile Asp Gly Tyr His Val Pro Ala Gly Thr Thr Ala Met Val  
 405 410 415

Asn Thr Trp Ala Ile Cys Arg Asp Pro His Val Trp Lys Asp Pro Leu  
 420 425 430

Glu Phe Met Pro Glu Arg Phe Val Thr Ala Gly Gly Asp Ala Glu Phe  
 435 440 445

Ser Ile Leu Gly Ser Asp Pro Arg Leu Ala Pro Phe Gly Ser Gly Arg  
 450 455 460

Arg Ala Cys Pro Gly Lys Thr Leu Gly Trp Ala Thr Val Asn Phe Trp  
 465 470 475 480

Val Ala Ser Leu Leu His Glu Phe Glu Trp Val Pro Ser Asp Glu Lys  
 485 490 495

Gly Val Asp Leu Thr Glu Val Leu Lys Leu Ser Ser Glu Met Ala Asn  
 500 505 510

Pro Leu Thr Val Lys Val Arg Pro Arg Arg Gly  
 515 520

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1788 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 6..1601

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGGTC ATG GGC ATG GCC ATG GAT GCT TTC CAG CAC CAA ACT CTC ATT	47
Met Gly Met Ala Met Asp Ala Phe Gln His Gln Thr Leu Ile	
1 5 10	
TCC ATC ATT CTG GCC ATG TTA GTA GGC GTG TTG ATT TAT GGC TTA AAG	95
Ser Ile Ile Leu Ala Met Leu Val Gly Val Leu Ile Tyr Gly Leu Lys	
15 20 25 30	
AGA ACA CAT AGT GGC CAT GGC AAG ATC TGT AGT GCA CCT CAA GCA GGA	143
Arg Thr His Ser Gly His Gly Lys Ile Cys Ser Ala Pro Gln Ala Gly	
35 40 45	

-58-

GGA GCA TGG CCA ATT ATT GGC CAT TTA CAC CTC TTT GGG GGT CAT CAA	191
Gly Ala Trp Pro Ile Ile Gly His Leu His Leu Phe Gly Gly His Gln	
50 55 60	
CAT ACT CAC AAA ACA CTT GGG ATA ATG GCA GAG AAA CAT GGA CCA ATT	239
His Thr His Lys Thr Leu Gly Ile Met Ala Glu Lys His Gly Pro Ile	
65 70 75	
TTC ACA ATA AAG CTT GGT TCA TAC AAA GTT CTT GTA TTG AGT AGC TGG	287
Phe Thr Ile Lys Leu Gly Ser Tyr Lys Val Leu Val Leu Ser Ser Trp	
80 85 90	
GAG ATG GCC AAG GAG TGT TTC ACT GTC CAT GAC AAA GCA TTT TCT ACC	335
Glu Met Ala Lys Glu Cys Phe Thr Val His Asp Lys Ala Phe Ser Thr	
95 100 105 110	
AGA CCC TGT GTT GCA GCC TCA AAG CTA ATG GGC TAC AAC TAT GCC ATG	383
Arg Pro Cys Val Ala Ala Ser Lys Leu Met Gly Tyr Asn Tyr Ala Met	
115 120 125	
TTT GGC TTC ACT CCT TAT GGT CCT TAT TGG CGT GAG ATA AGG AAA TTA	431
Phe Gly Phe Thr Pro Tyr Gly Pro Tyr Trp Arg Glu Ile Arg Lys Leu	
130 135 140	
ACT ACT ATT CAG CTT CTA TCT AAC CAC CGG CTT GAA CTG CTG AAG AAC	479
Thr Thr Ile Gln Leu Leu Ser Asn His Arg Leu Glu Leu Leu Lys Asn	
145 150 155	
ACA AGA ACA TCT GAG TCA GAA GTT GCA ATA AGA GAG CTT TAT AAG TTG	527
Thr Arg Thr Ser Glu Ser Glu Val Ala Ile Arg Glu Leu Tyr Lys Leu	
160 165 170	
TGG TCT AGA GAA GGT TGT CCA AAG GGA GGG GTT TTG GTA GAT ATG AAG	575
Trp Ser Arg Glu Gly Cys Pro Lys Gly Gly Val Leu Val Asp Met Lys	
175 180 185 190	
CAG TGG TTT GGG GAT TTA ACT CAT AAT ATT GTT CTG AGA ATG GTG AGA	623
Gln Trp Phe Gly Asp Leu Thr His Asn Ile Val Leu Arg Met Val Arg	
195 200 205	
GGG AAG CCA TAC TAT GAT GGT GCT AGT GAT GAT TAT GCA GAA GGT GAA	671
Gly Lys Pro Tyr Tyr Asp Gly Ala Ser Asp Asp Tyr Ala Glu Gly Glu	
210 215 220	
GCA AGA AGG TAC AAG AAA GTT ATG GGA GAG TGT GTG AGT TTG TTT GGG	719
Ala Arg Arg Tyr Lys Lys Val Met Gly Glu Cys Val Ser Leu Phe Gly	
225 230 235	
GTG TTT GTG TTA TCT GAT GCT ATT CCA TTT CTG GGG TGG TTG GAC ATC	767
Val Phe Val Leu Ser Asp Ala Ile Pro Phe Leu Gly Trp Leu Asp Ile	
240 245 250	
AAC GGA TAT GAA AAG GCC ATG AAG AGA ACT GCA AGT GAA TTG GAT CCT	815
Asn Gly Tyr Glu Lys Ala Met Lys Arg Thr Ala Ser Glu Leu Asp Pro	
255 260 265 270	
CTG GTT GAA GGG TGG TTA GAG GAA CAC AAA AGG AAA AGA GCT TTC AAT	863
Leu Val Glu Gly Trp Leu Glu Glu His Lys Arg Lys Arg Ala Phe Asn	

-59-

275	280	285	
ATG GAT GCA AAA GAA GAA CAG GAT AAT TTC ATG GAT GTC ATG CTG AAT Met Asp Ala Lys Glu Glu Gln Asp Asn Phe Met Asp Val Met Leu Asn 290 295 300			911
GTT CTG AAA GAT GCA GAG ATT TCT GGT TAT GAT TCA GAT ACC ATC ATC Val Leu Lys Asp Ala Glu Ile Ser Gly Tyr Asp Ser Asp Thr Ile Ile 305 310 315			959
AAG GCT ACT TGT CTG AAT CTG ATT TTA GCA GGA AGC GAC ACC ACC ATG Lys Ala Thr Cys Leu Asn Leu Ile Leu Ala Gly Ser Asp Thr Thr Met 320 325 330			1007
ATT TCA CTA ACA TGG GTG CTA TCT CTG CTA CTT AAC CAT CAA ATG GAA Ile Ser Leu Thr Trp Val Leu Ser Leu Leu Leu Asn His Gln Met Glu 335 340 345 350			1055
CTA AAA AAA GTC CAA GAT GAA TTG GAC ACT TAT ATT GGG AAG GAC AGG Leu Lys Lys Val Gln Asp Glu Leu Asp Thr Tyr Ile Gly Lys Asp Arg 355 360 365			1103
AAG GTG GAA GAA TCT GAC ATA ACC AAG TTG GTG TAC CTC CAA GCC ATT Lys Val Glu Glu Ser Asp Ile Thr Lys Leu Val Tyr Leu Gln Ala Ile 370 375 380			1151
GTG AAG GAA ACA ATG CGG CTG TAT CCA CCA AGT CCT CTT ATC ACC CTT Val Lys Glu Thr Met Arg Leu Tyr Pro Pro Ser Pro Leu Ile Thr Leu 385 390 395			1199
CGT GCA GCC ATG GAA GAC TGC ACC TTC TCA GGT GGC TAT CAC ATT CCT Arg Ala Ala Met Glu Asp Cys Thr Phe Ser Gly Gly Tyr His Ile Pro 400 405 410			1247
GCT GGG ACA CGT TTA ATG GTG AAT GCT TGG AAG ATC CAC CGG GAT GGT Ala Gly Thr Arg Leu Met Val Asn Ala Trp Lys Ile His Arg Asp Gly 415 420 425 430			1295
CGT GTT TGG AGT GAT CCT CAT GAT TTC AAG CCT GGA AGG TTC TTG ACA Arg Val Trp Ser Asp Pro His Asp Phe Lys Pro Gly Arg Phe Leu Thr 435 440 445			1343
AGC CAC AAA GAT GTT GAT GTG AAG GGT CAG AAC TAT GAG CTC GTC CCT Ser His Lys Asp Val Asp Val Lys Gly Gln Asn Tyr Glu Leu Val Pro 450 455 460			1391
TTT GGT TCT GGA AGG AGA GCA TGC CCT GGA GCC TCG CTG GCT CTG CGT Phe Gly Ser Gly Arg Arg Ala Cys Pro Gly Ala Ser Leu Ala Leu Arg 465 470 475			1439
GTG GTG CAC TTG ACC ATG GCT AGA CTG TTA CAT TCT TTC AAT GTT GCT Val Val His Leu Thr Met Ala Arg Leu Leu His Ser Phe Asn Val Ala 480 485 490			1487
TCT CCT TCA AAT CAA GTT GTG GAC ATG ACA GAG AGC ATT GGA CTC ACA Ser Pro Ser Asn Gln Val Val Asp Met Thr Glu Ser Ile Gly Leu Thr 495 500 505 510			1535
AAT TTA AAA GCA ACC CCG CTT GAA ATT CTC CTA ACT CCA CGT CTA GAC			1583

-60-

Asn Leu Lys Ala Thr Pro Leu Glu Ile Leu Leu Thr Pro Arg Leu Asp  
 515 520 525

ACC AAA CTT TAT GAG AAC TAGATTAAAT TAAGCTAGTT TTCTCCCAAA 1631  
 Thr Lys Leu Tyr Glu Asn  
 530

TAAGGGGAGG GGTCTCTAG GTCCTGAAAT CGGGTAATAA CAATAACATG GTTAATGCAG 1691

CTTCCATGTA GGATAATGAT TATTCACCTCA TGGGTCACCT TTTAATGGAG CCTCAGTGTA 1751

TTATAATAAC TCCAAACTTG TGGGTCACAA TCCCCC 1788

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 532 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Gly Met Ala Met Asp Ala Phe Gln His Gln Thr Leu Ile Ser Ile  
 1 5 10 15

Ile Leu Ala Met Leu Val Gly Val Leu Ile Tyr Gly Leu Lys Arg Thr  
 20 25 30

His Ser Gly His Gly Lys Ile Cys Ser Ala Pro Gln Ala Gly Gly Ala  
 35 40 45

Trp Pro Ile Ile Gly His Leu His Leu Phe Gly Gly His Gln His Thr  
 50 55 60

His Lys Thr Leu Gly Ile Met Ala Glu Lys His Gly Pro Ile Phe Thr  
 65 70 75 80

Ile Lys Leu Gly Ser Tyr Lys Val Leu Val Leu Ser Ser Trp Glu Met  
 85 90 95

Ala Lys Glu Cys Phe Thr Val His Asp Lys Ala Phe Ser Thr Arg Pro  
 100 105 110

Cys Val Ala Ala Ser Lys Leu Met Gly Tyr Asn Tyr Ala Met Phe Gly  
 115 120 125

Phe Thr Pro Tyr Gly Pro Tyr Trp Arg Glu Ile Arg Lys Leu Thr Thr  
 130 135 140

Ile Gln Leu Leu Ser Asn His Arg Leu Glu Leu Leu Lys Asn Thr Arg  
 145 150 155 160

Thr Ser Glu Ser Glu Val Ala Ile Arg Glu Leu Tyr Lys Leu Trp Ser  
 165 170 175

Arg Glu Gly Cys Pro Lys Gly Gly Val Leu Val Asp Met Lys Gln Trp  
 180 185 190

-61-

Phe Gly Asp Leu Thr His Asn Ile Val Leu Arg Met Val Arg Gly Lys  
 195 200 205  
 Pro Tyr Tyr Asp Gly Ala Ser Asp Asp Tyr Ala Glu Gly Glu Ala Arg  
 210 215 220  
 Arg Tyr Lys Lys Val Met Gly Glu Cys Val Ser Leu Phe Gly Val Phe  
 225 230 235 240  
 Val Leu Ser Asp Ala Ile Pro Phe Leu Gly Trp Leu Asp Ile Asn Gly  
 245 250 255  
 Tyr Glu Lys Ala Met Lys Arg Thr Ala Ser Glu Leu Asp Pro Leu Val  
 260 265 270  
 Glu Gly Trp Leu Glu Glu His Lys Arg Lys Arg Ala Phe Asn Met Asp  
 275 280 285  
 Ala Lys Glu Glu Gln Asp Asn Phe Met Asp Val Met Leu Asn Val Leu  
 290 295 300  
 Lys Asp Ala Glu Ile Ser Gly Tyr Asp Ser Asp Thr Ile Ile Lys Ala  
 305 310 315 320  
 Thr Cys Leu Asn Leu Ile Leu Ala Gly Ser Asp Thr Thr Met Ile Ser  
 325 330 335  
 Leu Thr Trp Val Leu Ser Leu Leu Leu Asn His Gln Met Glu Leu Lys  
 340 345 350  
 Lys Val Gln Asp Glu Leu Asp Thr Tyr Ile Gly Lys Asp Arg Lys Val  
 355 360 365  
 Glu Glu Ser Asp Ile Thr Lys Leu Val Tyr Leu Gln Ala Ile Val Lys  
 370 375 380  
 Glu Thr Met Arg Leu Tyr Pro Pro Ser Pro Leu Ile Thr Leu Arg Ala  
 385 390 395 400  
 Ala Met Glu Asp Cys Thr Phe Ser Gly Gly Tyr His Ile Pro Ala Gly  
 405 410 415  
 Thr Arg Leu Met Val Asn Ala Trp Lys Ile His Arg Asp Gly Arg Val  
 420 425 430  
 Trp Ser Asp Pro His Asp Phe Lys Pro Gly Arg Phe Leu Thr Ser His  
 435 440 445  
 Lys Asp Val Asp Val Lys Gly Gln Asn Tyr Glu Leu Val Pro Phe Gly  
 450 455 460  
 Ser Gly Arg Arg Ala Cys Pro Gly Ala Ser Leu Ala Leu Arg Val Val  
 465 470 475 480  
 His Leu Thr Met Ala Arg Leu Leu His Ser Phe Asn Val Ala Ser Pro  
 485 490 495  
 Ser Asn Gln Val Val Asp Met Thr Glu Ser Ile Gly Leu Thr Asn Leu  
 500 505 510

-62-

Lys Ala Thr Pro Leu Glu Ile Leu Leu Thr Pro Arg Leu Asp Thr Lys  
           515                                  520                                  525

Leu Tyr Glu Asn  
       530

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1657 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1548

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CTT GTT CTT CTT TCT CTA TTG TCT ATA GTC ATC TCC ATT GTT CTC TTC	48
Leu Val Leu Leu Ser Leu Leu Ser Ile Val Ile Ser Ile Val Leu Phe	
1                                  5                                  10                                  15	
ATT ACC CAC ACA CAC AAA AGA AAC AAC ACT CCA AGA GGA CCA CCA GGT	96
Ile Thr His Thr His Lys Arg Asn Asn Thr Pro Arg Gly Pro Pro Gly	
20                                  25                                  30	
CCT CCA CCT CTT CCT CTC ATC GGC AAC CTT CAC CAA CTC CAC AAC TCA	144
Pro Pro Pro Leu Pro Leu Ile Gly Asn Leu His Gln Leu His Asn Ser	
35                                  40                                  45	
TCC CCA CAT CTC TGC CTA TGG CAA CTC GCC AAA CTC CAC GGT CCT CTC	192
Ser Pro His Leu Cys Leu Trp Gln Leu Ala Lys Leu His Gly Pro Leu	
50                                  55                                  60	
ATG TCG TTT CGC CTC GGC GCC GTG CAA ACC GTC GTG GTT TCA TCG GCC	240
Met Ser Phe Arg Leu Gly Ala Val Gln Thr Val Val Val Ser Ser Ala	
65                                  70                                  75                                  80	
AGA ATC GCC GAA CAA ATC TTG AAA ACC CAC GAC CTC AAC TTC GCT TCC	288
Arg Ile Ala Glu Gln Ile Leu Lys Thr His Asp Leu Asn Phe Ala Ser	
85                                  90                                  95	
AGG CCT CTC TTC GTG GGC CCG AGA AAG CTC TCT TAC GAC GGG TTG GAC	336
Arg Pro Leu Phe Val Gly Pro Arg Lys Leu Ser Tyr Asp Gly Leu Asp	
100                                  105                                  110	
ATG GGC TTC GCA CCG TAC GGC CCG TAC TGG AGA GAA ATG AAG AAA CTC	384
Met Gly Phe Ala Pro Tyr Gly Pro Tyr Trp Arg Glu Met Lys Lys Leu	
115                                  120                                  125	
TGC ATC GTT CAC CTC TTC AGC GCG CAA CGC GTT CGG TCC TTT CGA CCA	432
Cys Ile Val His Leu Phe Ser Ala Gln Arg Val Arg Ser Phe Arg Pro	



-63-

130	135	140	
ATT CGA GAG AAC GAG GTT GCA AAA ATG GTT CGG AAA CTG TCG GAA CAC Ile Arg Glu Asn Glu Val Ala Lys Met Val Arg Lys Leu Ser Glu His 145 150 155 160			480
GAA GCT TCG GGT ACT GTC GTG AAC TTG ACC GAA ACT TTG ATG TCT TTC Glu Ala Ser Gly Thr Val Val Asn Leu Thr Glu Thr Leu Met Ser Phe 165 170 175			528
ACG AAC TCT TTG ATA TGC AGA ATC GCG TTG GGG AAA AGT TAC GGT TGT Thr Asn Ser Leu Ile Cys Arg Ile Ala Leu Gly Lys Ser Tyr Gly Cys 180 185 190			576
GAG TAC GAG GAA GTA GTT GTT GAT GAG GTA CTG GGA AAC CGG AGG AGC Glu Tyr Glu Glu Val Val Val Asp Glu Val Leu Gly Asn Arg Arg Ser 195 200 205			624
AGG TTG CAG GTT CTG CTC AAC GAG GCT CAA GCG TTG CTT TCG GAG TTT Arg Leu Gln Val Leu Leu Asn Glu Ala Gln Ala Leu Leu Ser Glu Phe 210 215 220			672
TTC TTT TCG GAT TAT TTT CCG CCT ATA GGA AAG TGG GTT GAT AGA GTG Phe Phe Ser Asp Tyr Phe Pro Pro Ile Gly Lys Trp Val Asp Arg Val 225 230 235 240			720
ACG GGA ATT CTA TCG CGG CTT GAT AAA ACG TTC AAG GAG TTG GAC GCG Thr Gly Ile Leu Ser Arg Leu Asp Lys Thr Phe Lys Glu Leu Asp Ala 245 250 255			768
TGC TAC GAA CGA TCA TCC TAT GAT CAC ATG GAT TCG GCA AAG AGT GGT Cys Tyr Glu Arg Ser Ser Tyr Asp His Met Asp Ser Ala Lys Ser Gly 260 265 270			816
AAA AAA GAT AAT GAC AAC AAA GAA GTC AAA GAT ATT ATT GAT ATT CTT Lys Lys Asp Asn Asp Asn Lys Glu Val Lys Asp Ile Ile Asp Ile Leu 275 280 285			864
CTC CAG CTA CTT GAT GAT CGT TCC TTC ACC TTT GAT CTC ACT CTC GAC Leu Gln Leu Leu Asp Asp Arg Ser Phe Thr Phe Asp Leu Thr Leu Asp 290 295 300			912
CAC ATA AAA GCC GTG CTC ATG AAC ATC TTT ATA GCA GGA ACA GAC CCG His Ile Lys Ala Val Leu Met Asn Ile Phe Ile Ala Gly Thr Asp Pro 305 310 315 320			960
AGT TCC GCG ACA ATA GTT TGG GCA ATG AAT GCA CTG TTG AAG AAT CCC Ser Ser Ala Thr Ile Val Trp Ala Met Asn Ala Leu Leu Lys Asn Pro 325 330 335			1008
AAT GTG ATG AGC AAG GTT CAA GGA GAA GTG AGA AAT CTA TTC GGT GAC Asn Val Met Ser Lys Val Gln Gly Glu Val Arg Asn Leu Phe Gly Asp 340 345 350			1056
AAA GAT TTC ATA AAC GAA GAT GAT GTC GAA AGC CTT CCT TAT CTC AAA Lys Asp Phe Ile Asn Glu Asp Asp Val Glu Ser Leu Pro Tyr Leu Lys 355 360 365			1104
GCA GTG GTG AAG GAG ACA TTA AGA TTA TTC CCA CCT TCA CCA CTA CTT			1152

-64-

Ala Val Val Lys Glu Thr Leu Arg Leu Phe Pro Pro Ser Pro Leu Leu	
370 375 380	
TTG CCA AGG GTA ACA ATG GAA ACA TGC AAC ATA GAA GGG TAC GAA ATT	1200
Leu Pro Arg Val Thr Met Glu Thr Cys Asn Ile Glu Gly Tyr Glu Ile	
385 390 395 400	
CAA GCC AAA ACT ATA GTG CAT GTT AAT GCA TGG GCC ATA GCA AGG GAC	1248
Gln Ala Lys Thr Ile Val His Val Asn Ala Trp Ala Ile Ala Arg Asp	
405 410 415	
CCT GAG AAT TGG GAA GAG CCT GAG AAA TTT TTC CCC GAA AGG TTC CTT	1296
Pro Glu Asn Trp Glu Glu Pro Glu Lys Phe Phe Pro Glu Arg Phe Leu	
420 425 430	
GAG AGT TCG ATG GAG TTA AAG GGG AAT GAT GAG TTT AAG GTG ATC CCG	1344
Glu Ser Ser Met Glu Leu Lys Gly Asn Asp Glu Phe Lys Val Ile Pro	
435 440 445	
TTT GGT TCT GGA AGG AGA ATG TGT CCT GCG AAG CAC ATG GGA ATT ATG	1392
Phe Gly Ser Gly Arg Arg Met Cys Pro Ala Lys His Met Gly Ile Met	
450 455 460	
AAT GTT GAG CTT TCT CTT GCT AAT CTC ATT CAC ACG TTT GAT TGG GAA	1440
Asn Val Glu Leu Ser Leu Ala Asn Leu Ile His Thr Phe Asp Trp Glu	
465 470 475 480	
GTG GCT AAA GGG TTC GAC AAG GAA GAA ATG TTG GAC ACG CAA ATG AAA	1488
Val Ala Lys Gly Phe Asp Lys Glu Glu Met Leu Asp Thr Gln Met Lys	
485 490 495	
CCA GGA ATA ACG ATG CAC AAG AAA AGT GAT CTT TAC CTA GTG GCA AAG	1536
Pro Gly Ile Thr Met His Lys Lys Ser Asp Leu Tyr Leu Val Ala Lys	
500 505 510	
AAA CCG ACA ACG TAGCACACGT TGGTACATTC ACTATAACAC ACAAGAAAGT	1588
Lys Pro Thr Thr	
515	
TGATAATGAC TTGTGTATGC AACTATGCTC TATGCACTAT GCACTATGTT TATTGACCAT	1648
TAATTACTG	1657

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 516 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Leu Val Leu Leu Ser Leu Leu Ser Ile Val Ile Ser Ile Val Leu Phe
1 5 10 15
Ile Thr His Thr His Lys Arg Asn Asn Thr Pro Arg Gly Pro Pro Gly
20 25 30

-65-

Pro Pro Pro Leu Pro Leu Ile Gly Asn Leu His Gln Leu His Asn Ser  
           35                          40                          45  
 Ser Pro His Leu Cys Leu Trp Gln Leu Ala Lys Leu His Gly Pro Leu  
           50                          55                          60  
 Met Ser Phe Arg Leu Gly Ala Val Gln Thr Val Val Val Ser Ser Ala  
           65                          70                          75                          80  
 Arg Ile Ala Glu Gln Ile Leu Lys Thr His Asp Leu Asn Phe Ala Ser  
                           85                          90                          95  
 Arg Pro Leu Phe Val Gly Pro Arg Lys Leu Ser Tyr Asp Gly Leu Asp  
                           100                          105                          110  
 Met Gly Phe Ala Pro Tyr Gly Pro Tyr Trp Arg Glu Met Lys Lys Leu  
           115                          120                          125  
 Cys Ile Val His Leu Phe Ser Ala Gln Arg Val Arg Ser Phe Arg Pro  
           130                          135                          140  
 Ile Arg Glu Asn Glu Val Ala Lys Met Val Arg Lys Leu Ser Glu His  
           145                          150                          155                          160  
 Glu Ala Ser Gly Thr Val Val Asn Leu Thr Glu Thr Leu Met Ser Phe  
                           165                          170                          175  
 Thr Asn Ser Leu Ile Cys Arg Ile Ala Leu Gly Lys Ser Tyr Gly Cys  
                           180                          185                          190  
 Glu Tyr Glu Glu Val Val Val Asp Glu Val Leu Gly Asn Arg Arg Ser  
           195                          200                          205  
 Arg Leu Gln Val Leu Leu Asn Glu Ala Gln Ala Leu Leu Ser Glu Phe  
           210                          215                          220  
 Phe Phe Ser Asp Tyr Phe Pro Pro Ile Gly Lys Trp Val Asp Arg Val  
           225                          230                          235                          240  
 Thr Gly Ile Leu Ser Arg Leu Asp Lys Thr Phe Lys Glu Leu Asp Ala  
                           245                          250                          255  
 Cys Tyr Glu Arg Ser Ser Tyr Asp His Met Asp Ser Ala Lys Ser Gly  
           260                          265                          270  
 Lys Lys Asp Asn Asp Asn Lys Glu Val Lys Asp Ile Ile Asp Ile Leu  
           275                          280                          285  
 Leu Gln Leu Leu Asp Asp Arg Ser Phe Thr Phe Asp Leu Thr Leu Asp  
           290                          295                          300  
 His Ile Lys Ala Val Leu Met Asn Ile Phe Ile Ala Gly Thr Asp Pro  
           305                          310                          315                          320  
 Ser Ser Ala Thr Ile Val Trp Ala Met Asn Ala Leu Leu Lys Asn Pro  
                           325                          330                          335  
 Asn Val Met Ser Lys Val Gln Gly Glu Val Arg Asn Leu Phe Gly Asp  
           340                          345                          350

-66-

Lys	Asp	Phe	Ile	Asn	Glu	Asp	Val	Glu	Ser	Leu	Pro	Tyr	Leu	Lys		
355						360						365				
Ala	Val	Val	Lys	Glu	Thr	Leu	Arg	Leu	Phe	Pro	Pro	Ser	Pro	Leu	Leu	
370						375						380				
Leu	Pro	Arg	Val	Thr	Met	Glu	Thr	Cys	Asn	Ile	Glu	Gly	Tyr	Glu	Ile	
385						390						395			400	
Gln	Ala	Lys	Thr	Ile	Val	His	Val	Asn	Ala	Trp	Ala	Ile	Ala	Arg	Asp	
			405						410						415	
Pro	Glu	Asn	Trp	Glu	Glu	Pro	Glu	Lys	Phe	Phe	Pro	Glu	Arg	Phe	Leu	
			420						425						430	
Glu	Ser	Ser	Met	Glu	Leu	Lys	Gly	Asn	Asp	Glu	Phe	Lys	Val	Ile	Pro	
435						440						445				
Phe	Gly	Ser	Gly	Arg	Arg	Met	Cys	Pro	Ala	Lys	His	Met	Gly	Ile	Met	
450						455						460				
Asn	Val	Glu	Leu	Ser	Leu	Ala	Asn	Leu	Ile	His	Thr	Phe	Asp	Trp	Glu	
465						470						475			480	
Val	Ala	Lys	Gly	Phe	Asp	Lys	Glu	Glu	Met	Leu	Asp	Thr	Gln	Met	Lys	
			485						490						495	
Pro	Gly	Ile	Thr	Met	His	Lys	Lys	Ser	Asp	Leu	Tyr	Leu	Val	Ala	Lys	
			500						505						510	
Lys	Pro	Thr	Thr													
515																

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1824 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS  
(B) LOCATION: 54..1616

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGAAAATTAG CCTCACAAA GCAAAGATCA AACAAACCAA GGACGAGAAC ACG ATG 56  
Met  
1

TTG CTT GAA CTT GCA CTT GGT TTA TTG GTT TTG GCT CTG TTT CTG CAC 104  
Leu Leu Glu Leu Ala Leu Gly Leu Val Leu Ala Leu Phe Leu His  
5 10 15

-67-

TTG CGT CCC ACA CCC ACT GCA AAA TCA AAA GCA CTT CGC CAT CTC CCA	152
Leu Arg Pro Thr Pro Thr Ala Lys Ser Lys Ala Leu Arg His Leu Pro	
20 25 30	
AAC CCA CCA AGC CCA AAG CCT CGT CTT CCC TTC ATA GGA CAC CTT CAT	200
Asn Pro Pro Ser Pro Lys Pro Arg Leu Pro Phe Ile Gly His Leu His	
35 40 45	
CTC TTA AAA GAC AAA CTT CTC CAC TAC GCA CTC ATC GAC CTC TCC AAA	248
Leu Leu Lys Asp Lys Leu Leu His Tyr Ala Leu Ile Asp Leu Ser Lys	
50 55 60 65	
AAA CAT GGT CCC TTA TTC TCT CTC TAC TTT GGC TCC ATG CCA ACC GTT	296
Lys His Gly Pro Leu Phe Ser Leu Tyr Phe Gly Ser Met Pro Thr Val	
70 75 80	
GTT GCC TCC ACA CCA GAA TTG TTC AAG CTC TTC CTC CAA ACG CAC GAG	344
Val Ala Ser Thr Pro Glu Leu Phe Lys Leu Phe Leu Gln Thr His Glu	
85 90 95	
GCA ACT TCC TTC AAC ACA AGG TTC CAA ACC TCA GCC ATA AGA CGC CTC	392
Ala Thr Ser Phe Asn Thr Arg Phe Gln Thr Ser Ala Ile Arg Arg Leu	
100 105 110	
ACC TAT GAT AGC TCA GTG GCC ATG GTT CCC TTC GGA CCT TAC TGG AAG	440
Thr Tyr Asp Ser Ser Val Ala Met Val Pro Phe Gly Pro Tyr Trp Lys	
115 120 125	
TTC GTG AGG AAG CTC ATC ATG AAC GAC CTT CCC AAC GCC ACC ACT GTA	488
Phe Val Arg Lys Leu Ile Met Asn Asp Leu Pro Asn Ala Thr Thr Val	
130 135 140 145	
AAC AAG TTG AGG CCT TTG AGG ACC CAA CAG ACC CGC AAG TTC CTT AGG	536
Asn Lys Leu Arg Pro Leu Arg Thr Gln Gln Thr Arg Lys Phe Leu Arg	
150 155 160	
GTT ATG GCC CAA GGC GCA GAG GCA CAG AAG CCC CTT GAC TTG ACC GAG	584
Val Met Ala Gln Gly Ala Glu Ala Gln Lys Pro Leu Asp Leu Thr Glu	
165 170 175	
GAG CTT CTG AAA TGG ACC AAC AGC ACC ATC TCC ATG ATG ATG CTC GGC	632
Glu Leu Leu Lys Trp Thr Asn Ser Thr Ile Ser Met Met Met Leu Gly	
180 185 190	
GAG GCT GAG GAG ATC AGA GAC ATC GCT CGC GAG GTT CTT AAG ATC TTT	680
Glu Ala Glu Glu Ile Arg Asp Ile Ala Arg Glu Val Leu Lys Ile Phe	
195 200 205	
GGC GAA TAC AGC CTC ACT GAC TTC ATC TGG CCA TTG AAG CAT CTC AAG	728
Gly Glu Tyr Ser Leu Thr Asp Phe Ile Trp Pro Leu Lys His Leu Lys	
210 215 220 225	
GTT GGA AAG TAT GAG AAG AGG ATC GAC GAC ATC TTG AAC AAG TTC GAC	776
Val Gly Lys Tyr Glu Lys Arg Ile Asp Asp Ile Leu Asn Lys Phe Asp	
230 235 240	
CCT GTC GTT GAA AGG GTC ATC AAG AAG CGC CGT GAG ATC GTG AGG AGG	824
Pro Val Val Glu Arg Val Ile Lys Lys Arg Arg Glu Ile Val Arg Arg	
245 250 255	

-68-

AGA AAG AAC GGA GAG GTT GTT GAG GGT GAG GTC AGC GGG GTT TTC CTT	872
Arg Lys Asn Gly Glu Val Val Glu Gly Glu Val Ser Gly Val Phe Leu	
260 265 270	
GAC ACT TTG CTT GAA TTC GCT GAG GAT GAG ACC ATG GAG ATC AAA ATC	920
Asp Thr Leu Leu Glu Phe Ala Glu Asp Glu Thr Met Glu Ile Lys Ile	
275 280 285	
ACC AAG GAC CAC ATC GAG GGT CTT GTT GTC GAC TTT TTC TCG GCA GGA	968
Thr Lys Asp His Ile Glu Gly Leu Val Val Asp Phe Phe Ser Ala Gly	
290 295 300 305	
ACA GAC TCC ACA GCG GTG GCA ACA GAG TGG GCA TTG GCA GAA CTC ATC	1016
Thr Asp Ser Thr Ala Val Ala Thr Glu Trp Ala Leu Ala Glu Leu Ile	
310 315 320	
AAC AAT CCT AAG GTG TTG GAA AAG GCT CGT GAG GAG GTC TAC AGT GTT	1064
Asn Asn Pro Lys Val Leu Glu Lys Ala Arg Glu Glu Val Tyr Ser Val	
325 330 335	
GTG GGA AAG GAC AGA CTT GTG GAC GAA GTT GAC ACT CAA AAC CTT CCT	1112
Val Gly Lys Asp Arg Leu Val Asp Glu Val Asp Thr Gln Asn Leu Pro	
340 345 350	
TAC ATT AGA GCA ATC GTG AAG GAG ACA TTC CGC ATG CAC CCG CCA CTC	1160
Tyr Ile Arg Ala Ile Val Lys Glu Thr Phe Arg Met His Pro Pro Leu	
355 360 365	
CCA GTG GTC AAA AGA AAG TGC ACA GAA GAG TGT GAG ATT AAT GGA TAT	1208
Pro Val Val Lys Arg Lys Cys Thr Glu Glu Cys Glu Ile Asn Gly Tyr	
370 375 380 385	
GTG ATC CCA GAG GGA GCA TTG ATT CTC TTC AAT GTA TGG CAA GTA GGA	1256
Val Ile Pro Glu Gly Ala Leu Ile Leu Phe Asn Val Trp Gln Val Gly	
390 395 400	
AGA GAC CCC AAA TAC TGG GAC AGA CCA TCG GAG TTC CGT CCT GAG AGG	1304
Arg Asp Pro Lys Tyr Trp Asp Arg Pro Ser Glu Phe Arg Pro Glu Arg	
405 410 415	
TTC CTA GAG ACA GGG GCT GAA GGG GAA GCA GGG CCT CTT GAT CTT AGG	1352
Phe Leu Glu Thr Gly Ala Glu Gly Glu Ala Gly Pro Leu Asp Leu Arg	
420 425 430	
GGA CAA CAT TTT CAA CTT CTC CCA TTT GGG TCT GGG AGG AGA ATG TGC	1400
Gly Gln His Phe Gln Leu Leu Pro Phe Gly Ser Gly Arg Arg Met Cys	
435 440 445	
CCT GGA GTC AAT CTG GCT ACT TCG GGA ATG GCA ACA CTT CTT GCA TCT	1448
Pro Gly Val Asn Leu Ala Thr Ser Gly Met Ala Thr Leu Leu Ala Ser	
450 455 460 465	
CTT ATT CAG TGC TTC GAC TTG CAA GTG CTG GGT CCA CAA GGA CAG ATA	1496
Leu Ile Gln Cys Phe Asp Leu Gln Val Leu Gly Pro Gln Gly Gln Ile	
470 475 480	
TTG AAG GGT GGT GAC GCC AAA GTT AGC ATG GAA GAG AGA GCC GGC CTC	1544
Leu Lys Gly Gly Asp Ala Lys Val Ser Met Glu Glu Arg Ala Gly Leu	
485 490 495	

-69-

ACT GTT CCA AGG GCA CAT AGT CTT GTC TGT GTT CCA CTT GCA AGG ATC 1592  
 Thr Val Pro Arg Ala His Ser Leu Val Cys Val Pro Leu Ala Arg Ile  
 500 505 510

GGC GTT GCA TCT AAA CTC CTT TCT TAATTAAGAT CATCATCATA TATAATATTT 1646  
 Gly Val Ala Ser Lys Leu Leu Ser  
 515 520

ACTTTTGTG TGTGATAAT CATCATTTCA ATAAGGTCTC GTTCATCTAC TTTTATGAA 1706  
 GTATATAAGC CCTTCCATGC ACATTGTATC ATCTCCCATTT TGTCTTCGTT TGCTACCTAA 1766  
 GGCAATCTTT TTTTTTTATAG AATCACATCA TCCTACTATA AACTATCAAT CCTTATAT 1824

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 521 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Leu Leu Glu Leu Ala Leu Gly Leu Leu Val Leu Ala Leu Phe Leu  
 1 5 10 15

His Leu Arg Pro Thr Pro Thr Ala Lys Ser Lys Ala Leu Arg His Leu  
 20 25 30

Pro Asn Pro Pro Ser Pro Lys Pro Arg Leu Pro Phe Ile Gly His Leu  
 35 40 45

His Leu Leu Lys Asp Lys Leu Leu His Tyr Ala Leu Ile Asp Leu Ser  
 50 55 60

Lys Lys His Gly Pro Leu Phe Ser Leu Tyr Phe Gly Ser Met Pro Thr  
 65 70 75 80

Val Val Ala Ser Thr Pro Glu Leu Phe Lys Leu Phe Leu Gln Thr His  
 85 90 95

Glu Ala Thr Ser Phe Asn Thr Arg Phe Gln Thr Ser Ala Ile Arg Arg  
 100 105 110

Leu Thr Tyr Asp Ser Ser Val Ala Met Val Pro Phe Gly Pro Tyr Trp  
 115 120 125

Lys Phe Val Arg Lys Leu Ile Met Asn Asp Leu Pro Asn Ala Thr Thr  
 130 135 140

Val Asn Lys Leu Arg Pro Leu Arg Thr Gln Gln Thr Arg Lys Phe Leu  
 145 150 155 160

Arg Val Met Ala Gln Gly Ala Glu Ala Gln Lys Pro Leu Asp Leu Thr  
 165 170 175

Glu Glu Leu Leu Lys Trp Thr Asn Ser Thr Ile Ser Met Met Met Leu

180

185

190

Gly Glu Ala Glu Glu Ile Arg Asp Ile Ala Arg Glu Val Leu Lys Ile  
195 200 205

Phe Gly Glu Tyr Ser Leu Thr Asp Phe Ile Trp Pro Leu Lys His Leu  
210 215 220

Lys Val Gly Lys Tyr Glu Lys Arg Ile Asp Asp Ile Leu Asn Lys Phe  
225 230 235 240

Asp Pro Val Val Glu Arg Val Ile Lys Lys Arg Arg Glu Ile Val Arg  
245 250 255

Arg Arg Lys Asn Gly Glu Val Val Glu Gly Glu Val Ser Gly Val Phe  
260 265 270

Leu Asp Thr Leu Leu Glu Phe Ala Glu Asp Glu Thr Met Glu Ile Lys  
275 280 285

Ile Thr Lys Asp His Ile Glu Gly Leu Val Val Asp Phe Phe Ser Ala  
290 295 300

Gly Thr Asp Ser Thr Ala Val Ala Thr Glu Trp Ala Leu Ala Glu Leu  
305 310 315 320

Ile Asn Asn Pro Lys Val Leu Glu Lys Ala Arg Glu Glu Val Tyr Ser  
325 330 335

Val	Val	Gly	Lys	Asp	Arg	Leu	Val	Asp	Glu	Val	Asp	Thr	Gln	Asn	Leu
			340					345					350		

Pro Tyr Ile Arg Ala Ile Val Lys Glu Thr Phe Arg Met His Pro Pro  
355 360 365

Leu Pro Val Val Lys Arg Lys Cys Thr Glu Glu Cys Glu Ile Asn Gly  
370 375 380

Tyr Val Ile Pro Glu Gly Ala Leu Ile Leu Phe Asn Val Trp Gln Val  
385                      390                      395                      400

Gly Arg Asp Pro Lys Tyr Trp Asp Arg Pro Ser Glu Phe Arg Pro Glu  
405 410 415

Arg Phe Leu Glu Thr Gly Ala Glu Gly Glu Ala Gly Pro Leu Asp Leu  
420 425 430

Arg Gly Gln His Phe Gln Leu Leu Pro Phe Gly Ser Gly Arg Arg Met  
435 440 445

Cys Pro Gly Val Asn Leu Ala Thr Ser Gly Met Ala Thr Leu Leu Ala  
450 455 460

Ser Leu Ile Gln Cys Phe Asp Leu Gln Val Leu Gly Pro Gln Gly Gln  
465 470 475 480

Ile Leu Lys Gly Gly Asp Ala Lys Val Ser Met Glu Glu Arg Ala Gly  
485 490 495

Leu Thr Val Pro Arg Ala His Ser Leu Val Cys Val Pro Leu Ala Arg



-71-

500 505 510

Ile Gly Val Ala Ser Lys Leu Leu Ser  
515 520

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1831 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 20..1747

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CAACACTCGC AGTACCGCC ATG AGT GTC GAC ACT TCC TCC ACC CTC TCC ACC	52
Met Ser Val Asp Thr Ser Ser Thr Leu Ser Thr	
1 5 10	
GTC ACC GAT GCC AAT CTT CAC TCC AGA TTT CAT TCT CGT CTT GTT CCA	100
Val Thr Asp Ala Asn Leu His Ser Arg Phe His Ser Arg Leu Val Pro	
15 20 25	
TTC ACT CAT CAT TTC TCA CTT TCT CAA CCC AAA CGG ATT TCT TCA ATC	148
Phe Thr His His Phe Ser Leu Ser Gln Pro Lys Arg Ile Ser Ser Ile	
30 35 40	
AGA TGC CAA TCA ATT AAT ACC GAT AAG AAG AAA TCA AGT AGA AAT CTG	196
Arg Cys Gln Ser Ile Asn Thr Asp Lys Lys Lys Ser Ser Arg Asn Leu	
45 50 55	
CTG GGC AAT GCA AGT AAC CTC CTC ACG GAC TTA TTA AGT GGT GGA AGT	244
Leu Gly Asn Ala Ser Asn Leu Leu Thr Asp Leu Leu Ser Gly Gly Ser	
60 65 70 75	
ATA GGG TCT ATG CCC ATA GCT GAA GGT GCA GTC TCA GAT CTG CTT GGT	292
Ile Gly Ser Met Pro Ile Ala Glu Gly Ala Val Ser Asp Leu Leu Gly	
80 85 90	
CGA CCT CTC TTT TTC TCA CTG TAT GAT TGG TTC TTG GAG CAT GGT GCG	340
Arg Pro Leu Phe Phe Ser Leu Tyr Asp Trp Phe Leu Glu His Gly Ala	
95 100 105	
GTG TAT AAA CTT GCC TTT GGA CCA AAA GCA TTT GTT GTT GTA TCA GAT	388
Val Tyr Lys Leu Ala Phe Gly Pro Lys Ala Phe Val Val Val Ser Asp	
110 115 120	
CCC ATA GTT GCT AGA CAT ATT CTG CGA GAA AAT GCA TTT TCT TAT GAC	436
Pro Ile Val Ala Arg His Ile Leu Arg Glu Asn Ala Phe Ser Tyr Asp	
125 130 135	
AAG GGA GTA CTT GCT GAT ATC CTT GAA CCA ATA ATG GGC AAA GGA CTC	484

-72-

Lys Gly Val Leu Ala Asp Ile Leu Glu Pro Ile Met Gly Lys Gly Leu 140 145 150 155	
ATA CCA GCA GAC CTT GAT ACT TGG AAG CAA AGG AGA AGA GTC ATT GCT Ile Pro Ala Asp Leu Asp Thr Trp Lys Gln Arg Arg Arg Val Ile Ala 160 165 170	532
CCG GCT TTC CAT AAC TCA TAC TTG GAA GCT ATG GTT AAA ATA TTC ACA Pro Ala Phe His Asn Ser Tyr Leu Glu Ala Met Val Lys Ile Phe Thr 175 180 185	580
ACT TGT TCA GAA AGA ACA ATA TTG AAG TTT AAT AAG CTT CTT GAA GGA Thr Cys Ser Glu Arg Thr Ile Leu Lys Phe Asn Lys Leu Leu Glu Gly 190 195 200	628
GAG GGT TAT GAT GGA CCT GAC TCA ATT GAA TTG GAT CTT GAG GCA GAG Glu Gly Tyr Asp Gly Pro Asp Ser Ile Glu Leu Asp Leu Glu Ala Glu 205 210 215	676
TTT TCT AGT TTG GCT CTT GAT ATT ATT GGG CTT GGT GTG TTC AAC TAT Phe Ser Ser Leu Ala Leu Asp Ile Ile Gly Leu Gly Val Phe Asn Tyr 220 225 230 235	724
GAC TTT GGT TCT GTC ACC AAA GAA TCT CCA GTT ATT AAG GCA GTC TAT Asp Phe Gly Ser Val Thr Lys Glu Ser Pro Val Ile Lys Ala Val Tyr 240 245 250	772
GGC ACT CTT TTT GAA GCT GAA CAC AGA TCC ACT TTC TAC ATT CCA TAT Gly Thr Leu Phe Glu Ala Glu His Arg Ser Thr Phe Tyr Ile Pro Tyr 255 260 265	820
TGG AAA ATT CCA TTG GCA AGG TGG ATA GTC CCA AGG CAA AGA AAG TTT Trp Lys Ile Pro Leu Ala Arg Trp Ile Val Pro Arg Gln Arg Lys Phe 270 275 280	868
CAG GAT GAC CTA AAG GTC ATC AAT ACT TGT CTT GAT GGA CTT ATC AGA Gln Asp Asp Leu Lys Val Ile Asn Thr Cys Leu Asp Gly Leu Ile Arg 285 290 295	916
AAT GCA AAA GAG AGC AGA CAG GAA ACA GAT GTT GAG AAA TTG CAG CAG Asn Ala Lys Glu Ser Arg Gln Glu Thr Asp Val Glu Lys Leu Gln Gln 300 305 310 315	964
AGG GAT TAC TTA AAT TTG AAG GAT GCA AGT CTT CTG CGT TTC CTG GTT Arg Asp Tyr Leu Asn Leu Lys Asp Ala Ser Leu Leu Arg Phe Leu Val 320 325 330	1012
GAT ATG CGG GGA GCT GAT GTT GAT GAT CGT CAG TTG AGG GAT GAT TTA Asp Met Arg Gly Ala Asp Val Asp Asp Arg Gln Leu Arg Asp Asp Leu 335 340 345	1060
ATG ACA ATG CTT ATT GCC GGT CAT GAA ACA ACG GCT GCA GTT CTT ACT Met Thr Met Leu Ile Ala Gly His Glu Thr Thr Ala Ala Val Leu Thr 350 355 360	1108
TGG GCA GTT TTC CTC CTA GCT CAA AAT CCT AGC AAA ATG AAG AAG GCT Trp Ala Val Phe Leu Leu Ala Gln Asn Pro Ser Lys Met Lys Lys Ala 365 370 375	1156

-73-

CAA GCA GAG GTA GAT TTG GTG CTG GGT ACG GGG AGG CCA ACT TTT GAA	1204
Gln Ala Glu Val Asp Leu Val Leu Gly Thr Gly Arg Pro Thr Phe Glu	
380 385 390 395	
TCA CTT AAG GAA TTG CAG TAC ATT AGA TTG ATT GTT GTG GAG GCT CTT	1252
Ser Leu Lys Glu Leu Gln Tyr Ile Arg Leu Ile Val Val Glu Ala Leu	
400 405 410	
CGT TTA TAC CCC CAA CCA CCT TTG CTG ATT AGA CGT TCA CTC AAA TCT	1300
Arg Leu Tyr Pro Gln Pro Pro Leu Leu Ile Arg Arg Ser Leu Lys Ser	
415 420 425	
GAT GTT TTA CCA GGT GGG CAC AAA GGT GAA AAA GAT GGT TAT GCA ATT	1348
Asp Val Leu Pro Gly Gly His Lys Gly Glu Lys Asp Gly Tyr Ala Ile	
430 435 440	
CCT GCT GGG ACT GAT GTC TTC ATT TCT GTA TAT AAT CTC CAT AGA TCT	1396
Pro Ala Gly Thr Asp Val Phe Ile Ser Val Tyr Asn Leu His Arg Ser	
445 450 455	
CCA TAT TTT TGG GAC CGC CCT GAT GAC TTC GAA CCA GAG AGA TTT CTT	1444
Pro Tyr Phe Trp Asp Arg Pro Asp Asp Phe Glu Pro Glu Arg Phe Leu	
460 465 470 475	
GTG CAA AAC AAG AAT GAA GAA ATT GAA GGA TGG GCT GGT CTT GAT CCA	1492
Val Gln Asn Lys Asn Glu Glu Ile Glu Gly Trp Ala Gly Leu Asp Pro	
480 485 490	
TCT CGA AGT CCC GGA GCC TTG TAT CCG AAC GAG GTT ATA TCG GAT TTT	1540
Ser Arg Ser Pro Gly Ala Leu Tyr Pro Asn Glu Val Ile Ser Asp Phe	
495 500 505	
GCA TTC TTA CCT TTT GGT GGC GGA CCA CGA AAA TGT GTT GGG GAC CAA	1588
Ala Phe Leu Pro Phe Gly Gly Pro Arg Lys Cys Val Gly Asp Gln	
510 515 520	
TTT GCT CTG ATG GAG TCC ACT GTA GCG TTG ACT ATG CTG CTC CAG AAT	1636
Phe Ala Leu Met Glu Ser Thr Val Ala Leu Thr Met Leu Leu Gln Asn	
525 530 535	
TTT GAC GTG GAA CTA AAA GGG ACC CCT GAA TCG GTG GAA CTA GTT ACT	1684
Phe Asp Val Glu Leu Lys Gly Thr Pro Glu Ser Val Glu Leu Val Thr	
540 545 550 555	
GGG GCA ACT ATT CAT ACC AAA AAT GGA ATG TGG TGC AGA TTG AAG AAG	1732
Gly Ala Thr Ile His Thr Lys Asn Gly Met Trp Cys Arg Leu Lys Lys	
560 565 570	
AGA TCT AAT TTA CGT TGACATATGT ACTGTGGCCA TTTTCTTAT ACAGAATAAT	1787
Arg Ser Asn Leu Arg	
575	
GTATATTATT ATTCTTTGAG AATAATATGA ATAAATTCCT AGAC	1831

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 576 amino acids

-74-

(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

```

Met Ser Val Asp Thr Ser Ser Thr Leu Ser Thr Val Thr Asp Ala Asn
 1             5             10             15
Leu His Ser Arg Phe His Ser Arg Leu Val Pro Phe Thr His His Phe
          20             25             30
Ser Leu Ser Gln Pro Lys Arg Ile Ser Ser Ile Arg Cys Gln Ser Ile
          35             40             45
Asn Thr Asp Lys Lys Lys Ser Ser Arg Asn Leu Leu Gly Asn Ala Ser
          50             55             60
Asn Leu Leu Thr Asp Leu Leu Ser Gly Gly Ser Ile Gly Ser Met Pro
          65             70             75             80
Ile Ala Glu Gly Ala Val Ser Asp Leu Leu Gly Arg Pro Leu Phe Phe
          85             90             95
Ser Leu Tyr Asp Trp Phe Leu Glu His Gly Ala Val Tyr Lys Leu Ala
          100            105            110
Phe Gly Pro Lys Ala Phe Val Val Val Ser Asp Pro Ile Val Ala Arg
          115            120            125
His Ile Leu Arg Glu Asn Ala Phe Ser Tyr Asp Lys Gly Val Leu Ala
          130            135            140
Asp Ile Leu Glu Pro Ile Met Gly Lys Gly Leu Ile Pro Ala Asp Leu
          145            150            155            160
Asp Thr Trp Lys Gln Arg Arg Arg Val Ile Ala Pro Ala Phe His Asn
          165            170            175
Ser Tyr Leu Glu Ala Met Val Lys Ile Phe Thr Thr Cys Ser Glu Arg
          180            185            190
Thr Ile Leu Lys Phe Asn Lys Leu Leu Glu Gly Glu Gly Tyr Asp Gly
          195            200            205
Pro Asp Ser Ile Glu Leu Asp Leu Glu Ala Glu Phe Ser Ser Leu Ala
          210            215            220
Leu Asp Ile Ile Gly Leu Gly Val Phe Asn Tyr Asp Phe Gly Ser Val
          225            230            235            240
Thr Lys Glu Ser Pro Val Ile Lys Ala Val Tyr Gly Thr Leu Phe Glu
          245            250            255
Ala Glu His Arg Ser Thr Phe Tyr Ile Pro Tyr Trp Lys Ile Pro Leu
          260            265            270
Ala Arg Trp Ile Val Pro Arg Gln Arg Lys Phe Gln Asp Asp Leu Lys
          275            280            285

```

-75-

Val Ile Asn Thr Cys Leu Asp Gly Leu Ile Arg Asn Ala Lys Glu Ser  
 290 295 300  
 Arg Gln Glu Thr Asp Val Glu Lys Leu Gln Gln Arg Asp Tyr Leu Asn  
 305 310 315 320  
 Leu Lys Asp Ala Ser Leu Leu Arg Phe Leu Val Asp Met Arg Gly Ala  
 325 330 335  
 Asp Val Asp Asp Arg Gln Leu Arg Asp Asp Leu Met Thr Met Leu Ile  
 340 345 350  
 Ala Gly His Glu Thr Thr Ala Ala Val Leu Thr Trp Ala Val Phe Leu  
 355 360 365  
 Leu Ala Gln Asn Pro Ser Lys Met Lys Lys Ala Gln Ala Glu Val Asp  
 370 375 380  
 Leu Val Leu Gly Thr Gly Arg Pro Thr Phe Glu Ser Leu Lys Glu Leu  
 385 390 395 400  
 Gln Tyr Ile Arg Leu Ile Val Val Glu Ala Leu Arg Leu Tyr Pro Gln  
 405 410 415  
 Pro Pro Leu Leu Ile Arg Arg Ser Leu Lys Ser Asp Val Leu Pro Gly  
 420 425 430  
 Gly His Lys Gly Glu Lys Asp Gly Tyr Ala Ile Pro Ala Gly Thr Asp  
 435 440 445  
 Val Phe Ile Ser Val Tyr Asn Leu His Arg Ser Pro Tyr Phe Trp Asp  
 450 455 460  
 Arg Pro Asp Asp Phe Glu Pro Glu Arg Phe Leu Val Gln Asn Lys Asn  
 465 470 475 480  
 Glu Glu Ile Glu Gly Trp Ala Gly Leu Asp Pro Ser Arg Ser Pro Gly  
 485 490 495  
 Ala Leu Tyr Pro Asn Glu Val Ile Ser Asp Phe Ala Phe Leu Pro Phe  
 500 505 510  
 Gly Gly Gly Pro Arg Lys Cys Val Gly Asp Gln Phe Ala Leu Met Glu  
 515 520 525  
 Ser Thr Val Ala Leu Thr Met Leu Leu Gln Asn Phe Asp Val Glu Leu  
 530 535 540  
 Lys Gly Thr Pro Glu Ser Val Glu Leu Val Thr Gly Ala Thr Ile His  
 545 550 555 560  
 Thr Lys Asn Gly Met Trp Cys Arg Leu Lys Lys Arg Ser Asn Leu Arg  
 565 570 575

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1704 base pairs

-76-

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS  
 (B) LOCATION: 38..1564

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CAGGCTCCAC AAAACATCTC ATCATTACACC CAACAAA ATG GCG CTG CTT CTG ATA	55
Met Ala Leu Leu Leu Ile	
1 5	
ATT CCC ATC TCA CTG GTC ACC CTC TGG CTC GGT TAC ACC CTA TAC CAG	103
Ile Pro Ile Ser Leu Val Thr Leu Trp Leu Gly Tyr Thr Leu Tyr Gln	
10 15 20	
CGA TTA AGA TTC AAG CTC CCT CCG GGT CCA CGG CCC TGG CCG GTA GTC	151
Arg Leu Arg Phe Lys Leu Pro Pro Gly Pro Arg Pro Trp Pro Val Val	
25 30 35	
GGT AAC CTC TAC GAC ATA AAA CCC GTC CGC TTC CGG TGC TTC GCG GAG	199
Gly Asn Leu Tyr Asp Ile Lys Pro Val Arg Phe Arg Cys Phe Ala Glu	
40 45 50	
TGG GCG CAG TCT TAC GGC CCC ATA ATA TCG GTT TGG TTC GGT TCG ACC	247
Trp Ala Gln Ser Tyr Gly Pro Ile Ile Ser Val Trp Phe Gly Ser Thr	
55 60 65 70	
CTA AAC GTC ATC GTT TCG AAC TCG GAG CTG GCG AAG GAG GTG CTG AAG	295
Leu Asn Val Ile Val Ser Asn Ser Glu Leu Ala Lys Glu Val Leu Lys	
75 80 85	
GAG CAC GAT CAG CTG CTG GCG GAC CGC CAC CGG AGC CGG TCG GCG GCG	343
Glu His Asp Gln Leu Leu Ala Asp Arg His Arg Ser Arg Ser Ala Ala	
90 95 100	
AAG TTC AGC CGC GAC GGG AAG GAT CTA ATT TGG GCC GAT TAT GGG CCG	391
Lys Phe Ser Arg Asp Gly Lys Asp Leu Ile Trp Ala Asp Tyr Gly Pro	
105 110 115	
CAC TAC GTG AAG GTG AGG AAG GTT TGC ACG CTC GAG CTT TTC TCG CCG	439
His Tyr Val Lys Val Arg Lys Val Cys Thr Leu Glu Leu Phe Ser Pro	
120 125 130	
AAG CGC CTC GAG GCC CTG AGG CCC ATT AGG GAG GAC GAG GTC ACC TCC	487
Lys Arg Leu Glu Ala Leu Arg Pro Ile Arg Glu Asp Glu Val Thr Ser	
135 140 145 150	
ATG GTT GAC TCC GTT TAC AAT CAC TGC ACC AGC ACT GAA AAT TTG GGG	535
Met Val Asp Ser Val Tyr Asn His Cys Thr Ser Thr Glu Asn Leu Gly	
155 160 165	
AAA GGA ATA TTG TTG AGG AAG CAC TTG GGG GTT GTG GCA TTC AAC AAC	583
Lys Gly Ile Leu Leu Arg Lys His Leu Gly Val Val Ala Phe Asn Asn	

-77-

170	175	180	
ATA ACC AGG TTG GCA TTT GGG AAA AGA TTT GTG AAC TCA GAA GGT GTG Ile Thr Arg Leu Ala Phe Gly Lys Arg Phe Val Asn Ser Glu Gly Val 185 190 195			631
ATG GAT GAG CAA GGA GTA GAA TTC AAG GCC ATT GTG GAA AAT GGG TTA Met Asp Glu Gln Gly Val Glu Phe Lys Ala Ile Val Glu Asn Gly Leu 200 205 210			679
AAG CTA GGA GCA TCT CTA GCC ATG GCA GAA CAC ATC CCT TGG CTT CGC Lys Leu Gly Ala Ser Leu Ala Met Ala Glu His Ile Pro Trp Leu Arg 215 220 225 230			727
TGG ATG TTC CCA CTG GAA GAA GGA GCT TTT GCC AAG CAT GGA GCC CGC Trp Met Phe Pro Leu Glu Glu Gly Ala Phe Ala Lys His Gly Ala Arg 235 240 245			775
CGC GAC CGA CTC ACC AGA GCC ATC ATG GCA GAG CAC ACT GAA GCA CGC Arg Asp Arg Leu Thr Arg Ala Ile Met Ala Glu His Thr Glu Ala Arg 250 255 260			823
AAG AAA TCT GGT GGT GCC AAG CAA CAT TTT GTT GAT GCC CTC CTC ACA Lys Lys Ser Gly Gly Ala Lys Gln His Phe Val Asp Ala Leu Leu Thr 265 270 275			871
TTG CAA GAC AAA TAT GAC CTT AGT GAA GAC ACC ATC ATT GGT CTC CTT Leu Gln Asp Lys Tyr Asp Leu Ser Glu Asp Thr Ile Ile Gly Leu Leu 280 285 290			919
TGG GAT ATG ATC ACA GCA GGG ATG GAC ACA ACT GCA ATT TCA GTT GAG Trp Asp Met Ile Thr Ala Gly Met Asp Thr Thr Ala Ile Ser Val Glu 295 300 305 310			967
TGG GCC ATG GCT GAG TTG ATA AGA AAC CCA AGG GTG CAA CAA AAG GTC Trp Ala Met Ala Glu Leu Ile Arg Asn Pro Arg Val Gln Gln Lys Val 315 320 325			1015
CAA GAG GAG CTA GAC AGG GTA ATT GGG CTT GAA AGG GTG ATG ACT GAA Gln Glu Glu Leu Asp Arg Val Ile Gly Leu Glu Arg Val Met Thr Glu 330 335 340			1063
GCA GAC TTC TCA AAT CTC CCT TAC CTA CAA TGT GTG ACC AAA GAA GCA Ala Asp Phe Ser Asn Leu Pro Tyr Leu Gln Cys Val Thr Lys Glu Ala 345 350 355			1111
ATG AGG CTT CAC CCA CCA ACC CCA CTA ATG CTC CCA CAC CGT GCC AAT Met Arg Leu His Pro Pro Thr Pro Leu Met Leu Pro His Arg Ala Asn 360 365 370			1159
GCC AAT GTC AAA GTT GGA GGC TAT GAC ATT CCC AAA GGG TCC AAT GTG Ala Asn Val Lys Val Gly Gly Tyr Asp Ile Pro Lys Gly Ser Asn Val 375 380 385 390			1207
CAT GTG AAT GTG TGG GCG GTG GCC CGC GAC CCG GCC GTG TGG AAG GAT His Val Asn Val Trp Ala Val Ala Arg Asp Pro Ala Val Trp Lys Asp 395 400 405			1255
CCA TTG GAG TTC CGA CCC GAA AGG TTC CTT GAG GAG GAT GTA GAC ATG Pro Leu Glu Phe Arg Pro Glu Arg Phe Leu Glu Glu Asp Val Asp Met			1303

-78-

410	415	420	
AAG GGC CAT GAC TTT AGG CTA CTT CCA TTC GGG TCG GGT CGA CGA GTA			1351
Lys Gly His Asp Phe Arg Leu Leu Pro Phe Gly Ser Gly Arg Arg Val			
425	430	435	
TGC CCG GGT GCC CAA CTT GGT ATC AAC TTG GCA GCA TCC ATG TTG GGC			1399
Cys Pro Gly Ala Gln Leu Gly Ile Asn Leu Ala Ala Ser Met Leu Gly			
440	445	450	
CAC CTC TTG CAC CAT TTC TGT TGG ACC CCA CCT GAA GGA ATG AAG CCT			1447
His Leu Leu His His Phe Cys Trp Thr Pro Pro Glu Gly Met Lys Pro			
455	460	465	470
GAG GAA ATT GAC ATG GGA GAG AAT CCA GGG CTA GTC ACA TAC ATG AGG			1495
Glu Glu Ile Asp Met Gly Glu Asn Pro Gly Leu Val Thr Tyr Met Arg			
475	480	485	
ACT CCA ATA CAA GCT GTG GTT TCT CCT AGG CTC CCC TCA CAT TTA TAC			1543
Thr Pro Ile Gln Ala Val Val Ser Pro Arg Leu Pro Ser His Leu Tyr			
490	495	500	
AAA CGT GTG CCT GCT GAG ATC TAATCTTTCT TTTCTTTCCC TTGGACTACT			1594
Lys Arg Val Pro Ala Glu Ile			
505			
CTTTGTTGCA TTAAGAAAAA TGCCTTGTTGG CACTACTTTT ATCTTTGTGT TTATGTAAC			1654
ACATATGAAA TCACAATTTA AGGAACTAAG GAAAAACTCA TTGCGAGGGT			1704

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 509 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met	Ala	Leu	Leu	Leu	Ile	Ile	Pro	Ile	Ser	Leu	Val	Thr	Leu	Trp	Leu
1					5				10					15	
Gly	Tyr	Thr	Leu	Tyr	Gln	Arg	Leu	Arg	Phe	Lys	Leu	Pro	Pro	Gly	Pro
			20					25						30	
Arg	Pro	Trp	Pro	Val	Val	Gly	Asn	Leu	Tyr	Asp	Ile	Lys	Pro	Val	Arg
			35					40					45		
Phe	Arg	Cys	Phe	Ala	Glu	Trp	Ala	Gln	Ser	Tyr	Gly	Pro	Ile	Ile	Ser
		50					55					60			
Val	Trp	Phe	Gly	Ser	Thr	Leu	Asn	Val	Ile	Val	Ser	Asn	Ser	Glu	Leu
	65					70				75				80	
Ala	Lys	Glu	Val	Leu	Lys	Glu	His	Asp	Gln	Leu	Leu	Ala	Asp	Arg	His
				85					90					95	



-79-

Arg Ser Arg Ser Ala Ala Lys Phe Ser Arg Asp Gly Lys Asp Leu Ile  
 100 105 110  
 Trp Ala Asp Tyr Gly Pro His Tyr Val Lys Val Arg Lys Val Cys Thr  
 115 120 125  
 Leu Glu Leu Phe Ser Pro Lys Arg Leu Glu Ala Leu Arg Pro Ile Arg  
 130 135 140  
 Glu Asp Glu Val Thr Ser Met Val Asp Ser Val Tyr Asn His Cys Thr  
 145 150 155 160  
 Ser Thr Glu Asn Leu Gly Lys Gly Ile Leu Leu Arg Lys His Leu Gly  
 165 170 175  
 Val Val Ala Phe Asn Asn Ile Thr Arg Leu Ala Phe Gly Lys Arg Phe  
 180 185 190  
 Val Asn Ser Glu Gly Val Met Asp Glu Gln Gly Val Glu Phe Lys Ala  
 195 200 205  
 Ile Val Glu Asn Gly Leu Lys Leu Gly Ala Ser Leu Ala Met Ala Glu  
 210 215 220  
 His Ile Pro Trp Leu Arg Trp Met Phe Pro Leu Glu Glu Gly Ala Phe  
 225 230 235 240  
 Ala Lys His Gly Ala Arg Arg Asp Arg Leu Thr Arg Ala Ile Met Ala  
 245 250 255  
 Glu His Thr Glu Ala Arg Lys Lys Ser Gly Gly Ala Lys Gln His Phe  
 260 265 270  
 Val Asp Ala Leu Leu Thr Leu Gln Asp Lys Tyr Asp Leu Ser Glu Asp  
 275 280 285  
 Thr Ile Ile Gly Leu Leu Trp Asp Met Ile Thr Ala Gly Met Asp Thr  
 290 295 300  
 Thr Ala Ile Ser Val Glu Trp Ala Met Ala Glu Leu Ile Arg Asn Pro  
 305 310 315 320  
 Arg Val Gln Gln Lys Val Gln Glu Glu Leu Asp Arg Val Ile Gly Leu  
 325 330 335  
 Glu Arg Val Met Thr Glu Ala Asp Phe Ser Asn Leu Pro Tyr Leu Gln  
 340 345 350  
 Cys Val Thr Lys Glu Ala Met Arg Leu His Pro Pro Thr Pro Leu Met  
 355 360 365  
 Leu Pro His Arg Ala Asn Ala Asn Val Lys Val Gly Gly Tyr Asp Ile  
 370 375 380  
 Pro Lys Gly Ser Asn Val His Val Asn Val Trp Ala Val Ala Arg Asp  
 385 390 395 400  
 Pro Ala Val Trp Lys Asp Pro Leu Glu Phe Arg Pro Glu Arg Phe Leu  
 405 410 415

-80-

Glu Glu Asp Val Asp Met Lys Gly His Asp Phe Arg Leu Leu Pro Phe  
                   420                                  425                                  430  
 Gly Ser Gly Arg Arg Val Cys Pro Gly Ala Gln Leu Gly Ile Asn Leu  
                   435                                  440                                  445  
 Ala Ala Ser Met Leu Gly His Leu Leu His His Phe Cys Trp Thr Pro  
                   450                                  455                                  460  
 Pro Glu Gly Met Lys Pro Glu Glu Ile Asp Met Gly Glu Asn Pro Gly  
                   465                                  470                                  475                                  480  
 Leu Val Thr Tyr Met Arg Thr Pro Ile Gln Ala Val Val Ser Pro Arg  
                                   485                                  490                                  495  
 Leu Pro Ser His Leu Tyr Lys Arg Val Pro Ala Glu Ile  
                                   500                                  505

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TGTCTAACTC CTTCTTTTC

20

## (2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 13 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Phe Leu Pro Phe Gly Xaa Gly Xaa Arg Xaa Cys Xaa Gly  
 1                                  5                                  10

## (2) INFORMATION FOR SEQ ID NO:21:

-81-

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Phe Xaa Xaa Gly Xaa Xaa Xaa Cys Xaa Gly  
1 5 10

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Xaa Cys Xaa Gly  
1

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEOUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Pro Glu Glu Phe Xaa Pro Glu Arg Phe  
1 5

**THAT WHICH IS CLAIMED IS:**

1. An isolated DNA molecule comprising a sequence selected from the group consisting of:

5 a) SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, and SEQ ID NO:17;

b) DNA sequences which encode an enzyme having a sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6:, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, and SEQ ID NO:18;

10 c) DNA sequences which have at least about 90% sequence identity to the DNA of (a) or (b) above and which encode a cytochrome P450 enzyme; and

d) DNA sequences which differ from the DNA of (a) or (c) above due to the degeneracy of the genetic code.

2. A peptide encoded by a DNA sequence of claim 1.

3. A cytochrome p450 enzyme having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6:, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, and SEQ ID NO:18.

4. An isolated DNA molecule comprising a sequence selected from the group consisting of:

5 a) SEQ ID NO:1;  
b) DNA sequences which encode an enzyme having SEQ ID NO:2,;

c) DNA sequences which have at least about 90% sequence identity to the DNA of (a) or (b) above and which encode a cytochrome P450 enzyme; and

10 d) DNA sequences which differ from the DNA of (a) or (c) above due to the degeneracy of the genetic code.

5. A peptide encoded by a DNA sequence of claim 4.

6. A cytochrome p450 peptide having SEQ ID NO:2.

7. A DNA construct comprising an expression cassette, which construct comprising in the 5' to 3' direction, a promoter operable in a plant cell and a DNA segment according to claim 1 positioned downstream from said promoter and operatively associated therewith.

8. A DNA construct according to claim 7, wherein said promoter is constitutively active in plant cells.

9. A DNA construct according to claim 7, wherein said promoter is the 35S promoter from Cauliflower Mosaic virus.

10. A DNA construct according to claim 7, said construct further comprising a plasmid.

11. A DNA construct according to claim 7 carried by a plant transformation vector.

12. A DNA construct according to claim 7 carried by an *Agrobacterium tumefaciens* plant transformation vector.

13. A plant cell containing a DNA construct according to claim 7.

14. A transgenic plant comprising plant cells according to claim 13.

15. A transgenic plant according to claim 14, wherein said plant is a monocot.

16. A transgenic plant according to claim 14, wherein said plant is a dicot.

17. A DNA construct comprising an expression cassette, which construct comprising in the 5' to 3' direction, a promoter operable in a plant cell, and a DNA segment encoding a peptide of SEQ ID NO:2 positioned downstream from said promoter and operatively associated therewith.

18. A DNA construct according to claim 17, wherein said promoter is constitutively active in plant cells.

19. A DNA construct according to claim 17, wherein said promoter is the 35S promoter from Cauliflower Mosaic virus.

20. A DNA construct according to claim 17, said construct further comprising a plasmid.

21. A DNA construct according to claim 17 carried by a plant transformation vector.

22. A DNA construct according to claim 17 carried by an *Agrobacterium tumefaciens* plant transformation vector.

23. A plant cell containing a DNA construct according to claim 17.

24. A transgenic plant comprising plant cells according to claim 23.

25. A transgenic plant according to claim 24, wherein said plant is a monocot.

26. A transgenic plant according to claim 24, wherein said plant is a dicot.

27. A method of making a transgenic plant cell having an increased ability to metabolize phenylurea compounds compared to an untransformed plant cell, said method comprising:

- a) providing a plant cell;
- 5        b) transforming said plant cell with an exogenous DNA construct comprising, in the 5' to 3' direction, a promoter operable in a plant cell and a DNA sequence encoding a peptide of SEQ ID NO:2, said DNA sequence operably linked to said promoter.

28. A method according to claim 27, wherein said plant cell is from a member of the Solanaceae family.

29. A method according to claim 27, wherein said promoter is the 35S promoter from Cauliflower Mosaic virus.

30. A method according to claim 27, wherein said transforming step is carried out by bombarding said plant cell with microparticles carrying said DNA construct.

31. A method according to claim 27 wherein said transforming step is carried out by infecting said plant cell with an *Agrobacterium tumefaciens* containing a Ti plasmid carrying said DNA construct.

32. A method according to claim 27, further comprising regenerating a plant from said transformed plant cell.

-86-

33. A transformed plant produced by the method of claim 32.

34. Seed or progeny of a plant according to claim 33, which seed or progeny has inherited said DNA sequence encoding a peptide of SEQ ID NO:2.

35. A transformed plant produced by the method of claim 32, which plant has increased resistance to phenylurea herbicides compared to wild-type plants of the same species.

36. A transgenic plant having an increased ability to metabolize phenylurea compounds compared to an untransformed plant cell, said transgenic plant comprising transgenic plant cells containing an exogenous DNA construct comprising, in the 5' to 3' direction, a promoter operable in said plant cell, said promoter operably linked to a DNA sequence encoding a peptide of SEQ ID NO:2.

37. A transgenic plant according to claim 36, wherein said promoter is the 35S promoter from Cauliflower Mosaic virus.

38. A transgenic plant according to claim 36, wherein said plant is a dicot.

39. A transgenic plant according to claim 36, wherein said plant is a monocot.

40. A transgenic plant according to claim 36, wherein said plant is a member of the family Solanaceae.

41. A transgenic plant according to claim 36, which plant is selected from the group consisting of tobacco, potato, tomato, corn, rice, cotton, soybean,



rape, wheat, oats, barley, rye and rice.

42. Progeny or seed of a plant according to claim 36, wherein said seed or progeny has inherited said DNA sequence encoding a peptide of SEQ ID NO:2.

43. A transformed plant according to claim 36, which plant has increased resistance to phenylurea herbicides compared to wild-type plants of the same species.

44. A crop comprising a plurality of plants according to claim 36 planted in an agricultural field.

45. A method of using a phenylurea herbicide as a post-emergence herbicide, comprising:

- a) planting a crop according to claim 44;
- b) applying to said crop a phenylurea herbicide.

46. A method according to claim 45, wherein said crop is selected from the group consisting of turfgrass, tobacco, potato, tomato, corn, rice, cotton, soybean, rape, wheat, oats, barley, rye and rice.

47. A method according to claim 45, wherein said herbicide is selected from the group consisting of fluometuron, linuron, chlortoluron and diuron.




Fig. 2




Fig. 1